TRANSACTIONS OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 00, Number 0, Pages 000–000
S 0002-9947(XX)0000-0

ON THE INVERSE SPECTRAL THEORY OF SCHRÖDINGER AND DIRAC OPERATORS

MIKLÓS HORVÁTH

ABSTRACT. We prove that under some conditions finitely many partially known spectra and partial information on the potential entirely determine the potential . This extends former results of Hochstadt, Lieberman, Gesztesy, Simon and others.

1. INTRODUCTION

Consider the Schrödinger operator

$$-y'' + q(x)y = \lambda y \quad \text{on} \quad (0,\pi) \tag{1.1}$$

with the boundary conditions

$$y(0)\cos\alpha + y'(0)\sin\alpha = 0 \tag{1.2}$$

$$y(\pi)\cos\beta + y'(\pi)\sin\beta = 0 \tag{1.3}$$

where $q \in L_1(0,\pi)$. The eigenvalues of (1.1)–(1.3) form the sequence

$$\lambda_0 < \lambda_1 < \dots .$$

Given a complex value λ define $y(x,\lambda)$ as the solution of (1.1) with the initial conditions

$$y(0) = \sin\alpha, \quad y'(0) = -\cos\alpha. \tag{1.4}$$

For $\lambda = \lambda_n$, $y(x,\lambda_n)$ is the eigenfunction of (1.1)–(1.3) corresponding to the eigenvalue λ_n. Introduce the normalizing constants

$$\alpha_n^2 = \int_0^\pi y^2(x,\lambda_n)\mathrm{d}x.$$

The *spectral function* of the problem (1.1)–(1.3) is defined to be

$$\varrho(\lambda) = \begin{cases} \sum\limits_{0<\lambda_n\leqslant\lambda} \frac{1}{\alpha_n^2} & \text{if} \quad \lambda > 0 \\ -\sum\limits_{\lambda<\lambda_n\leqslant 0} \frac{1}{\alpha_n^2} & \text{if} \quad \lambda < 0 \end{cases} \tag{1.5}$$

Received by the editors ????????
1991 *Mathematics Subject Classification.* Primary 34A55, 34B20; Secondary 34L40, 47A75.
Key words and phrases. Inverse spectral theory,m-function, spectral function.
Research supported by the Hungarian NSF Grant OTKA T#32374.

©1997 American Mathematical Society

A fundamental result of Marchenko [12] says that the spectral function determines the potential. A similar statement was given by G. Borg [2] on the *Titchmarsh-Weyl m-function* of the problem (1.1)–(1.3) as defined by

$$m(\lambda) = \frac{v'(0,\lambda)}{v(0,\lambda)} \qquad \lambda \in \mathbb{C}, \tag{1.6}$$

where $v(x,\lambda)$ is the solution of (1.1) with the initial conditions

$$v(\pi,\lambda) = \sin\beta, v'(\pi,\lambda) = -\cos\beta. \tag{1.7}$$

More precisely, let $q^*(x)$ be another potential and denote ϱ^* and m^* the corresponding functions. Then the two results together can be formulated as

Theorem 1.1. [2], [12]

a.) *If* $\varrho^*\ (\lambda) = \varrho(\lambda)$ *for* $\lambda \in \mathbb{R}$ *then* $q^*(x) = q(x)$ *a.e. in* $(0,\pi)$.
b.) *If* $m^*(\lambda) = m(\lambda)$ *for* $\lambda \in \mathbb{C} \setminus \mathbb{R}$ *then* $q^* = q$ *a.e. in* $(0,\pi)$.

The two statements are connected since m and ϱ can be expressed from each other (see e.g. [9]).
Now consider the eigenvalue problem corresponding to the Dirac operator

$$\begin{array}{rcl} u_2' + (V(x)+\mu)\,u_1 &=& \lambda u_1 \\ -u_1' + (V(x)-\mu)\,u_2 &=& \lambda u_2 \end{array} \qquad \text{on} \quad (0,\pi) \tag{1.8}$$

with the boundary conditions

$$u_1(0)\cos\alpha + u_2(0)\sin\alpha = 0 \tag{1.9}$$

$$u_1(\pi)\cos\beta + u_2(\pi)\sin\beta = 0. \tag{1.10}$$

Here we suppose that the potential $V(x)$ is continuous on $[0,\pi]$ and the mass μ is positive. The Dirac operator is the relativistic Schrödinger operator in quantum physics. The spectral function of the problem (1.8)–(1.10) is again defined by (1.5) with

$$\alpha_n^2 = \int_0^\pi \left[u_{n,1}^2(x,\lambda_n) + u_{n,2}^2(x,\lambda_n)\right] \mathrm{d}x, \qquad n \in \mathbb{Z},$$

where $(u_{n,1}(x,\lambda),\ u_{n,2}(x,\lambda))$ is the solution of (1.8) with the initial conditions

$$u_{n,1}(0,\lambda) = \sin\alpha,\ u_{n,2}(0,\lambda) = -\cos\alpha. \tag{1.11}$$

The relativistic version of Theorem 1.1

Theorem 1.2. [9], [13] *Let* ϱ^* *be the spectral function corresponding to the data* $V^*,\ \mu^*,\ \alpha^*$ *and* β^* *instead of* $V,\ \mu,\ \alpha$ *and* β. *If* $\varrho^* = \varrho$ *then* $V^* = V, \mu^* = \mu, \tan\alpha^* = \tan\alpha, \tan\beta^* = \tan\beta$.

Furthermore define the m-function as

$$m(\lambda) = \frac{v_2(0,\lambda)}{v_1(0,\lambda)}, \tag{1.12}$$

where $(v_1(x,\lambda),\ v_2(x,\lambda))$ is the solution of (1.8) with the initial conditions

$$v_1(\pi,\lambda) = \sin\beta,\ v_2(\pi,\lambda) = -\cos\beta. \tag{1.13}$$

Establishing the appropriate equation connecting ϱ and m we get

Theorem 1.3. *If* $m^*(\lambda) = m(\lambda)$ *then* $V^* = V,\ \mu^* = \mu$ *and* $\tan\beta^* = \tan\beta$.

Returning to the Schrödinger operator, it is known from Borg [1] that two spectra (defined by the boundary conditions (1.2)–(1.3) with α_1, β and α_2, β) determine the potential q. Hochstadt and Lieberman [7] proved that if half of the potential is known, then one spectrum is enough:

Theorem 1.4. [7] *Suppose that* $\cos\alpha \neq 0$ *and that* $q^* = q$ *a.e. on* $\left(0, \frac{\pi}{2}\right)$. *If the spectrum of (1.1)–(1.3) of* q *and of* q^* *is identical then* $q^* = q$ *a.e. on the whole* $(0, \pi)$.

Gesztesy and Simon [5] showed that more information on the potential can compensate for less information on the spectrum in the following sense:

Theorem 1.5. [5] *Let* $0 < \gamma < 1$ *and suppose that* $q^* = q$ *a.e. on* $\left(0, \frac{1+\gamma}{2}\pi\right)$, $\tan\beta^*$ *and* $\tan\beta \neq \infty$, $\tan\alpha^* = \tan\alpha \neq \infty$. *Denote* $S = \{\lambda_n : \lambda_n^* = \lambda_n\}$. *If*

$$\#\{\lambda \in S : \lambda \leqslant t_0\} \geqslant (1-\gamma)\#\{\lambda_n : \lambda_n < t_0\} + \alpha/2 \tag{1.14}$$

for all sufficiently large t_0 *then* $\tan\beta^* = \tan\beta$ *and* $q^* = q$ *a.e. on* $(0, \pi)$.

Later on del Rio, Gesztesy and Simon found the following related statements (see also [4]):

Theorem 1.6. [3] *Let* σ_N *and* σ_D *be the spectrum of (1.1)–(1.3) with* $\alpha = 0$, *respectively* $\alpha = \frac{\pi}{2}$. *Let* $S_N \subset \sigma_N$, $S_D \subset \sigma_D$ *and* $0 < \gamma < 1$. *If*

$$\#\{\lambda \in S_N \cup S_D : \lambda \leqslant t_0\} \geqslant (1-\gamma)\#\{\lambda \in \sigma_N \cup \sigma_D : \lambda \leqslant t_0\} \tag{1.15}$$

holds for all sufficiently large t_0, *then* S_N, S_D *and* q *on* $(0, \gamma\pi)$ *determine* q *a.e. on* $(0, \pi)$.

Theorem 1.7. [3] *Let* $\sigma_1, \sigma_2, \sigma_3$ *be the spectrum of (1.1)–(1.3) with* $\alpha_1, \alpha_2, \alpha_3$ *in (1.2). Let* $S_j \subset \sigma_j$ *satisfying*

$$\begin{aligned} &\#\{\lambda \in S_1 \cup S_2 \cup S_3 : \lambda \leqslant t_0\} \\ &\geqslant \frac{2}{3}\#\{\lambda \in \sigma_1 \cup \sigma_2 \cup \sigma_3 : \lambda \leqslant t_0\} \end{aligned} \tag{1.16}$$

for all sufficiently large t_0. *Then* S_1, S_2, S_3 *determine* q *a.e. on* $(0, \pi)$.

We will prove the following generalization of the above mentioned result of Borg and Theorems 1.4–1.7 (formulated on the segment $(0, \pi)$):

Theorem 1.8. *Let* σ_j, $j = 1, \dots, N$ *be the spectrum of (1.1)–(1.3) with* α_j *in (1.2). Suppose that* $\lambda_n^{(j)} \in \sigma_j$ *is known for* $n \in S_j$ *and denote*

$$n_j(t) = \sum_{\lambda_n^{(j)} < t^2,\ n \in S_j} 1$$

for $t \geqslant 0$. *Let* $0 \leqslant a < \pi$, $0 \leqslant \gamma \leqslant 1$, *and suppose that there exist* $t_0 > 0$ *and* $\delta > 0$ *such that for* $t \geqslant t_0$

$$\sum_{j=1}^{N} n_j(t) \geqslant \begin{cases} 2\left(1 - \frac{a}{\pi}\right)\left\{\gamma\left[t + \frac{1}{2}\right] + (1-\gamma)\left([t] + \frac{1}{2}\right)\right\} + \mathbf{O}\left(t^{-\delta}\right) \\ \qquad\qquad \text{if } \sin\beta \neq 0 \\ 2\left(1 - \frac{a}{\pi}\right)\left\{\gamma\left[t + \frac{1}{2}\right] + (1-\gamma)\left([t] + \frac{1}{2}\right)\right\} - 1 + \mathbf{O}\left(t^{-\delta}\right) \\ \qquad\qquad \text{if } \sin\beta = 0. \end{cases} \tag{1.17}$$

Then q on $(0,a)$ and the eigenvalues $\left\{\lambda_n^{(j)} : n \in S_j\right\}, j = 1,\dots,N$ determine q a.e. on $(0,\pi)$.

The counterpart for the Dirac case reads as follows:

Theorem 1.9. *Let $\sigma_j,\ j = 1\dots,N$ be the spectrum of (1.8)–(1.10) with α_j in (1.9). Suppose that $\lambda_n^{(j)} \in \sigma_j$ is known for $n \in S_j$ and denote*

$$\widetilde{n}_j(t) = \begin{cases} \sum\limits_{0<n<t,\ n\in S_j} 1 & \text{if} \quad t>0 \\ -\sum\limits_{t<n<0,\ n\in S_j} 1 & \text{if} \quad t<0. \end{cases}$$

(Thus, we will not count the value $n = 0$.) Suppose that the sets S_j are almost symmetrical, i.e. that $n \in S_j$ implies $-n \in S_j$ with finitely many possible exceptions and that the limits

$$\lim_{t\to\infty} \frac{\widetilde{n}_j(t)}{t} = \gamma_j \tag{1.18}$$

exist for $j = 1,\dots,N$. Denote d the number of indices j with $0 \in S_j$.
Let $\mu \in \mathbb{R}$, $0 \leqslant a < \pi$, $t_0 > 0$, $\delta > 0$ and $\varepsilon > 0$ be arbitrary numbers. Suppose finally that

$$\sum_{j=1}^{N} \widetilde{n}_j(t) \begin{cases} \geqslant 2\left(1-\frac{a}{\pi}\right)[t] - d + \mu - 1 + \varepsilon + \mathbf{O}\left(t^{-\delta}\right) \\ \qquad\qquad\qquad\qquad\qquad\qquad \text{if } t \geqslant t_0 \\ \leqslant -2\left(1-\frac{a}{\pi}\right)[-t] + \mu - 2\left(1-\frac{a}{\pi}\right) + \mathbf{O}\left(|t|^{-\delta}\right) \\ \qquad\qquad\qquad\qquad\qquad\qquad \text{if } t \leqslant -t_0. \end{cases} \tag{1.19}$$

Then V on $[0,a]$ and the eigenvalues $\left\{\lambda_n^{(j)} : n \in S_j\right\}$, $j = 1,...,N$ determine V on $[0,\pi]$.

2. Partial information on the potential

In this section we will prove Theorems 1.3, 1.8 and 1.9 . Though Theorem 1.3 might exist somewhere in the literature, I was not able to find it, so I will provide a proof. In order to do so, we need the Green function (or Green matrix) of the problem (1.8)–(1.10). It has the form

$$G(x,y,\lambda) = \text{p.v.} \sum_{-\infty}^{\infty} \frac{u(x,\lambda_n)u^T(y,\lambda_n)}{\alpha_n^2(\lambda-\lambda_n)}, \quad \lambda \neq \lambda_n. \tag{2.1}$$

where $u = \binom{u_1}{u_2}$, $\text{p.v.}\sum = \lim\limits_{N\to\infty} \sum\limits_{-N}^{N}$. The sum is convergent at every point (x,y) and the convergence is locally uniform outside the diagonal $x = y$. Consequently the Green matrix is continuous for $x < y$ and $x > y$, and the sum can be expressed in closed form (see e.g. Levitan, Sargsjan [10]):

$$G(x,y,\lambda) = \begin{cases} \frac{-1}{W(\lambda)} u(x,\lambda)v^T(y,\lambda) & \text{if } x<y \\ \frac{-1}{W(\lambda)} v(x,\lambda)u^T(y,\lambda) & \text{if } y<x. \end{cases} \tag{2.2}$$

Here $v = \binom{v_1}{v_2}$ and

$$W(\lambda) = u_1(x,\lambda)v_2(x,\lambda) - u_2(x,\lambda)v_1(x,\lambda).$$

This function W is independent of x since

$$\begin{aligned} W' =& u_1'v_2 - u_2'v_1 + u_1v_2' - u_2v_1' = (V-m-\lambda)u_2v_2 \\ & - (\lambda - V - m)u_1v_1 + (\lambda - V - m)u_1v_1 - (V-m-\lambda)u_2v_2 = 0. \end{aligned}$$

Lemma 2.1.

$$\begin{aligned} G_{11}(0,0,\lambda) &= \sin^2\alpha \cdot p.v. \int_{-\infty}^{\infty} \frac{d\rho(t)}{\lambda - t} \\ &= \sin\alpha\cos\alpha - \frac{\sin\alpha}{m(\lambda)\sin\alpha + \cos\alpha}, \end{aligned}$$

where G_{11} is the left upper element of the matrix G and p.v.$\int_{-\infty}^{\infty} = \lim_{R\to\infty}\int_{-R}^{R}$.

Proof. We are looking for the value of the jump

$$G_{11}(0,0+0,\lambda) - G_{11}(0,0,\lambda).$$

By definition

$$G_{11}(0,y,\lambda) = \sin\alpha \cdot \text{p.v.} \sum \frac{u_1(y,\lambda_n)}{\alpha_n^2(\lambda - \lambda_n)}.$$

Recall the asymptotical formulae

$$u_1(x,\lambda) = \sin\big(\lambda x - \int_0^x V + \alpha\big) + \mathbf{O}\big(\frac{e^{|\Im\lambda| x}}{|\lambda|}\big) \tag{2.3}$$

$$u_2(x,\lambda) = -\cos\left(\lambda x - \int_0^x V + \alpha\right) + \mathbf{O}\left(\frac{e^{|\Im\lambda| x}}{|\lambda|}\right), \quad |\lambda| \to \infty \tag{2.4}$$

with remainder terms uniform in x and

$$\alpha_n^2 = \pi + \mathbf{O}\big(\frac{1}{1+|n|}\big),$$

see [9]. Since $\alpha_n^2(\lambda - \lambda_n) = -\pi n + \mathbf{O}(1)$, we see that G_{11} has the same jump as

$$\sin\alpha \cdot \text{p.v.} \sum_{n\neq 0} \frac{\sin\big(\lambda_n y - \int_0^y V + \alpha\big)}{-\pi n} \tag{2.5}$$

(because the difference is continuous). Now consider the eigenvalue asymptotics:

$$\lambda_n = n + \frac{\vartheta}{\pi} + \mathbf{O}\left(\frac{1}{1+|n|}\right), \quad \vartheta = \beta - \alpha + \int_0^\pi V. \tag{2.6}$$

We get

$$\begin{aligned} \sin\big(\lambda_n y - \int_0^y V + \alpha\big) &= \sin\left(ny + a(y) + \mathbf{O}\Big(\frac{1}{1+|n|}\Big)\right), \\ a(y) &= \frac{\vartheta}{\pi} y - \int_0^y V + \alpha. \end{aligned}$$

So (2.5) has the same jump as

$$\sin\alpha \cdot \text{p.v.} \sum_{n\neq 0} \frac{\sin(ny+a(y))}{-\pi n} \tag{2.7}$$

$$= \sin\alpha \cdot \cos a(y) \cdot \frac{2}{-\pi} \sum_{n=1}^{\infty} \frac{\sin ny}{n}.$$

It is known that $\sum_{n=1}^{\infty} \frac{\sin ny}{n}$ is the Fourier series of $\frac{\pi-y}{2}$ on $(0,2\pi)$, extended 2π-periodically. From $y=0$ to $y=0+$ it has a jump $\frac{\pi}{2}$, so (2.7) has a jump $-\sin\alpha\cos\alpha$. Consequently by (2.5)

$$G_{11}(0,0,\lambda) - \sin\alpha\cos\alpha = \frac{-1}{W(\lambda)} u_1(0,\lambda)v_1(0,\lambda)$$

$$= -\frac{\sin\alpha\ v_1(0,\lambda)}{\sin\alpha\ v_2(0,\lambda) + \cos\alpha\ v_1(0,\lambda)} = \frac{-\sin\alpha}{m(\lambda)\sin\alpha + \cos\alpha}$$

as asserted. □

Lemma 2.2.

$$G_{22}(0,0,\lambda) = \cos^2\alpha \cdot \text{p.v.} \int_{-\infty}^{\infty} \frac{d\rho(t)}{\lambda - t}$$

$$= -\sin\alpha\cos\alpha + \frac{m(\lambda)\cos\alpha}{m(\lambda)\sin\alpha + \cos\alpha}.$$

Proof. Repeating the above reduction process we get that G_{22} has the same jump as the series

$$\cos\alpha \cdot \text{p.v.} \sum_{n\neq 0} \frac{\cos(ny+a(y))}{-\pi n} = \cos\alpha \sin a(y) \frac{2}{\pi} \sum_{1}^{\infty} \frac{\sin ny}{n}.$$

So this jump is $\sin\alpha\cos\alpha$ and then

$$G_{22}(0,0,\lambda) + \sin\alpha\cos\alpha = \frac{\cos\alpha\ v_2(0,\lambda)}{\sin\alpha\ v_2(0,\lambda) + \cos\alpha\ v_1(0,\lambda)}$$

$$= \frac{m(\lambda)\cos\alpha}{m(\lambda)\sin\alpha + \cos\alpha}$$

and we are done. □

Now the next statements follow immediately:

Corollary 2.3.

a) *If* $\sin\alpha \neq 0$, *then*

$$\text{p.v.} \int_{-\infty}^{\infty} \frac{d\rho(t)}{\lambda - t} = \cot\alpha - \frac{1}{\sin\alpha}\ \frac{1}{m(\lambda)\sin\alpha + \cos\alpha}$$

b) *If* $\cos\alpha \neq 0$, *then*

$$\text{p.v.} \int_{-\infty}^{\infty} \frac{d\rho(t)}{\lambda - t} = -\tan\alpha + \frac{1}{\cos\alpha}\ \frac{m(\lambda)}{m(\lambda)\sin\alpha + \cos\alpha}$$

c)

$$\text{p.v.}\int_{-\infty}^{\infty}\frac{d\rho(t)}{\lambda-t}=\frac{m(\lambda)\cos\alpha-\sin\alpha}{m(\lambda)\sin\alpha+\cos\alpha}.$$

Corollary 2.4. *The spectral function defines* $\tan\alpha$ *and the m-function, and conversely, the m-function and* $\tan\alpha$ *define the spectral function.*

Proof. The spectral function determines the mod π value of α, i.e. $\tan\alpha$, hence by Corollary 2.3 c) it defines also $m(\lambda)$. Conversely, $m(\lambda)$ and $\tan\alpha$ define the transform $\text{p.v.}\int_{-\infty}^{\infty}\frac{d\rho(t)}{\lambda-t}$ and then they define $\rho(t)$ itself by the Stieltjes inversion formula, see [10]. □

Proof. of Theorem 1.3 Define a boundary condition at $x=0$ with $\alpha=0$. Then by Corollary 2.3 b) $m(\lambda)=\text{p.v.}\int_{-\infty}^{\infty}\frac{d\rho(t)}{\lambda-t}$, and this defines ρ by the inversion formula. From ρ we determine V and $\tan\beta$. We are done. □

Next we analyze the growth properties of some infinite products in order to prove Theorems 1.8 and 1.9. Consider a sequence of arbitrary values

$$\lambda_n=n+\mathbf{O}(1)\quad n\in\mathbb{Z}. \tag{2.8}$$

Let S be a set of integers, almost symmetric with respect to the origin. This means that

$$S\subset\mathbb{Z},\quad n\in S\Rightarrow -n\in S. \tag{2.9}$$

with finitely many exceptions. Denote further

$$n_S(t)=\begin{cases}\displaystyle\sum_{0\leqslant\lambda_n\leqslant t}1 & \text{if } t>0\\ \displaystyle-\sum_{t<\lambda_n<0}1 & \text{if } t<0.\end{cases} \tag{2.10}$$

Lemma 2.5. *Suppose (2.8), (2.9) and that* $\lambda_n\neq 0$ *for* $n\in S$. *Then the product*

$$w_S(z)=\text{p.v.}\prod_{n\in S}\left(1-\frac{z}{\lambda_n}\right)$$

converges locally uniformly, defines an entire function with zeros λ_n. *Further we have*

$$\begin{aligned}\ln|w_S(z)| &= \text{p.v.}\int_{-\infty}^{\infty}\frac{n_S(t)}{t}\,\frac{|z|^2-xt}{|z|^2-2xt+t^2}\mathrm{d}t\\ &= \text{p.v.}\int_{-\infty}^{\infty}\frac{n_S(t)}{t}\,\frac{y^2-x(t-x)}{y^2+(t-x)^2}\mathrm{d}t.\end{aligned} \tag{2.11}$$

where $z=x+iy$.

Proof. The locally uniform convergence of the product follows from the simple estimate

$$\begin{aligned}\left(1-\frac{z}{\lambda_n}\right)\left(1-\frac{z}{\lambda_{-n}}\right)&=1-z\left(\frac{1}{\lambda_n}+\frac{1}{\lambda_{-n}}\right)+\frac{z^2}{\lambda_n\lambda_{-n}}\\ &=1+\mathbf{O}\left(\frac{1}{n^2}\right).\end{aligned}$$

To show (2.11) we integrate by parts

$$
\begin{aligned}
\ln|w_S(z)|^2 &= \text{p.v.}\sum \ln\left|1-\frac{z}{\lambda_n}\right|^2 \\
&= \text{p.v.}\sum \ln\left(\left(1-\frac{x}{\lambda_n}\right)^2+\frac{y^2}{\lambda_n^2}\right) \\
&= \text{p.v.}\sum \ln\left(1-\frac{2x}{\lambda_n}+\frac{|z|^2}{\lambda_n^2}\right) \\
&= \text{p.v.}\int_{-\infty}^{\infty} \ln\left(1-\frac{2x}{t}+\frac{|z|^2}{t^2}\right)\mathrm{d}n_S(t) \\
&= \text{p.v.}\left[\ln\left(1-\frac{2x}{t}+\frac{|z|^2}{t^2}\right)n_S(t)\right]_{-\infty}^{\infty} \\
&\quad -\text{p.v.}\int_{-\infty}^{\infty} n_S(t)\frac{\frac{2x}{t^2}-\frac{2|z|^2}{t^3}}{1-\frac{2x}{t}+\frac{|z|^2}{t^2}}\mathrm{d}t\,.
\end{aligned}
\tag{2.12}
$$

From

$$
\begin{aligned}
\ln\left(1-\frac{2x}{t}+\frac{|z|^2}{t^2}\right)n_S(t) &= \left(-\frac{2x}{t}+\frac{|z|^2}{t^2}\right)n_S(t)\big(1+\mathbf{o}(1)\big) \\
&= -2x\frac{n_S(t)}{t}+\mathbf{o}(1) \quad (t\to\pm\infty)
\end{aligned}
$$

the almost symmetrical property of S implies that

$$
\begin{aligned}
\ln|w_S(z)| &= \text{p.v.}\int_{-\infty}^{\infty} n_S(t)\frac{\frac{|z|^2}{t^3}-\frac{x}{t^2}}{1-\frac{2x}{t}+\frac{|z|^2}{t^2}}\mathrm{d}t \\
&= \text{p.v.}\int_{-\infty}^{\infty} \frac{n_S(t)}{t}\ \frac{|z|^2-xt}{t^2-2xt+|z|^2}\mathrm{d}t \\
&= \text{p.v.}\int_{-\infty}^{\infty} \frac{n_S(t)}{t}\ \frac{y^2-x(t-x)}{y^2+(t-x)^2}\mathrm{d}t.
\end{aligned}
$$

□

Lemma 2.6. *If we shift the zeros by* $\mathbf{O}\left(\frac{1}{n}\right)$*, the order of magnitude of* w_S *remains the same, except near the zeros. In other words, if*

$$
\lambda_n^* = \lambda_n + \mathbf{O}\left(\frac{1}{n}\right),\ \lambda_n^*\neq 0,\ w_S^*(z) = \text{p.v.}\prod_{n\in S}\left(1-\frac{z}{\lambda_n^*}\right)
$$

then for every $\delta>0$*,*

$$
|w_S^*(z)| \asymp |w_S(z)| \text{ if } |z-\lambda_n|>\delta,\ |z-\lambda_n^*|>\delta\ \forall n.
$$

The notation $\asymp$ *means that both* $\left|\frac{w_S^*(z)}{w_S(z)}\right|$ *and* $\left|\frac{w_S(z)}{w_S^*(z)}\right|$ *are bounded.*

Proof. We have to show that

$$
\ln|w_S^*(z)| - \ln|w_S(z)| \tag{2.13}
$$

is bounded whenever $|z-\lambda_n|>\delta$ and $|z-\lambda_n^*|>\delta$. For bounded z, $\ln|w_S^*(z)|$ and $\ln|w_S(z)|$ are both bounded, so we can suppose that $|z|$ is large, say, $|z|\geqslant K$. By the formula (2.11) we have

$$\ln|w_S^*(z)|-\ln|w_S(z)| = \text{p.v.}\int_{-\infty}^{\infty}\frac{n_S^*(t)-n_S(t)}{t}\,\frac{y^2-x(t-x)}{y^2+(t-x)^2}\mathrm{d}t. \tag{2.14}$$

Here $\frac{n_S^*(t)-n_S(t)}{t}$ takes values of order $\mathbf{O}\left(\frac{1}{n}\right)$ in segments of length $\mathbf{O}\left(\frac{1}{n}\right)$, where $t=\mathbf{O}(n)$, otherwise it is zero. It implies first that the integral in (2.14) is convergent also in the ordinary sense. Secondly, $\frac{y^2}{y^2+(t-x)^2}\leqslant 1$ implies that

$$\int_{-\infty}^{\infty}\frac{n_S^*(t)-n_S(t)}{t}\,\frac{y^2}{y^2+(t-x)^2}\mathrm{d}t=\mathbf{O}\left(\sum\frac{1}{n^2}\right)=\mathbf{O}(1) \tag{2.15}$$

uniformly in z (we used the fact that $n_S(t)$ and $n_S^*(t)$ are zero near $t=0$). So it remains to prove that

$$\int_{-\infty}^{\infty}\frac{n_S^*(t)-n_S(t)}{t}\,\frac{x(t-x)}{y^2+(t-x)^2}\mathrm{d}t=\mathbf{O}(1) \tag{2.16}$$
$$(|z|\geqslant K,\ |z-\lambda_n|>\delta,\ |z-\lambda_n^*|>\delta)$$

uniformly in z. If x is bounded then y is large, so

$$\frac{x(t-x)}{y^2+(t-x)^2}=\mathbf{O}\left(\frac{t-x}{1+(t-x)^2}\right)=\mathbf{O}(1)$$

and then an estimate of the type (2.15) applies. Hence we can suppose that x is large, e.g. $|x|\geqslant K/2$. Consider only $x>K/2$, since the case of negative x is similar. Split the integral (2.16) into four parts as follows. If $t\geqslant\frac{3x}{2}$ then $\frac{x(t-x)}{y^2+(t-x)^2}\asymp\frac{xt}{y^2+t^2}$, hence in (2.16)

$$\left|\int_{\frac{3x}{2}}^{\infty}\right|\leqslant c\sum_{k\geqslant\frac{3x}{2}}\frac{1}{k^2}\,\frac{xk}{y^2+k^2}\leqslant cx\sum_{k\geqslant\frac{3x}{2}}\frac{1}{k^2}\leqslant c$$

uniformly in $|z|$. If $\frac{x}{2}\leqslant t\leqslant\frac{3x}{2}$ then $y^2+(t-x)^2$ cannot be arbitrarily close to zero on the segments between λ_n^* and λ_n, hence

$$\frac{x(t-x)}{y^2+(t-x)^2}=\mathbf{O}\left(\frac{x|n-x|+x}{y^2+(n-x)^2+1}\right),\quad n=[t]$$

and then in (2.16)

$$\left|\int_{\frac{x}{2}}^{\frac{3x}{2}}\right|\leqslant c\sum_{k=1}^{\left[\frac{x}{2}\right]}\frac{1}{x^2}\,\frac{k\ x}{y^2+k^2}\leqslant c\sum_{k=1}^{\left[\frac{x}{2}\right]}\frac{1}{y^2+k^2}\leqslant c.$$

If $-\frac{x}{2}\leqslant t\leqslant\frac{x}{2}$, then $\frac{x(t-x)}{y^2+(t-x)^2}\asymp\frac{x^2}{y^2+x^2}=\mathbf{O}(1)$, hence $\left|\int_{-\frac{x}{2}}^{\frac{x}{2}}\right|\leqslant c$. Finally if $t\leqslant-\frac{x}{2}$, then $\frac{x(t-x)}{y^2+(t-x)^2}\asymp\frac{x\ t}{y^2+t^2}$, so in (2.16)

$$\left|\int_{\infty}^{-\frac{x}{2}}\right|\leqslant c\sum_{k=[\frac{x}{2}]}^{\infty}\frac{1}{k^2}\,\frac{x\ k}{y^2+k^2}\leqslant c\sum\frac{1}{y^2+k^2}\leqslant c.$$

This proves (2.16) and we are ready. □

Lemma 2.7. *Under the conditions of Lemma 2.5 we have for fixed x*

$$\ln|w_S(z)| = \int_{-\infty}^{\infty} \frac{n_S(t)}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t + \mathbf{O}(1) \quad (|y|\to\infty). \tag{2.17}$$

The $\mathbf{O}$*-term is locally uniform in* x*.*

Proof. We have to transform (2.11) into (2.17). First,

$$\begin{aligned}&\left|\int_{-\infty}^{\infty} \frac{n_S(t)}{t}\left[\frac{y^2}{y^2+(t-x)^2}-\frac{y^2}{y^2+t^2}\right]\mathrm{d}t\right|\\ \leqslant& c\int_{-\infty}^{\infty} y^2\frac{x^2+2|t|\ |x|}{(y^2+t^2)(y^2+(t-x)^2)}\mathrm{d}t\end{aligned}$$

and here

$$\begin{aligned}\int_{-\infty}^{\infty}\frac{y^2x^2}{(y^2+t^2)\,(y^2+(t-x)^2)}\mathrm{d}t &\leqslant x^2\int_{-\infty}^{\infty}\frac{\mathrm{d}t}{y^2+t^2}=\mathbf{O}(1),\\ \int_{|t|\leqslant|y|}\frac{y^2|t||x|}{\big(y^2+t^2\big)\big(y^2+(t-x)^2\big)}\mathrm{d}t &\leqslant |x|\int_{|t|\leqslant|y|}\frac{|t|}{y^2+t^2}\mathrm{d}t=\mathbf{O}(1),\\ \int_{|t|\geqslant|y|}\frac{y^2|t||x|}{\big(y^2+t^2\big)\big(y^2+(t-x)^2\big)}\mathrm{d}t &\leqslant cy^2|x|\int_{|t|\geqslant|y|}\frac{|t|}{t^4}\mathrm{d}t=\mathbf{O}(1).\end{aligned}$$

So it remains to show that

$$\text{p.v.}\int_{-\infty}^{\infty}\frac{n_S(t)}{t}\ \frac{t-x}{y^2+(t-x)^2}\mathrm{d}t=\mathbf{O}(1)\quad(|y|\to\infty) \tag{2.18}$$

locally uniformly in x. From (2.8) and (2.9) we see that

$$n_S(-t) = -n_S(t)+\mathbf{O}(t).$$

Taking into account that $\lambda_n\neq 0$, we get

$$\begin{aligned}&\text{p.v.}\int_{-\infty}^{\infty}\frac{n_S(t)}{t}\frac{t-x}{y^2+(t-x)^2}\mathrm{d}t\\ =&\int_{\delta}^{\infty}\frac{n_S(t)}{t}\left[\frac{t-x}{y^2+(t-x)^2}-\frac{t+x}{y^2+(t+x)^2}\right]\mathrm{d}t\\ &+\mathbf{O}\left(\int_{\delta}^{\infty}\frac{1}{t}\ \frac{|t+x|}{y^2+(t+x)^2}\mathrm{d}t\right).\end{aligned}$$

Now we have

$$\int_{\delta}^{\infty}\frac{1}{t}\ \frac{|t+x|}{y^2+(t+x)^2}\mathrm{d}t\leqslant c\int_{\delta}^{\infty}\frac{\mathrm{d}t}{y^2+(t+x)^2}=\mathbf{O}(1).$$

Furthermore

$$\begin{aligned}&\frac{t-x}{y^2+(t-x)^2}-\frac{t+x}{y^2+(t+x)^2}\\&=\frac{-2xy^2+(t-x)(t+x)2x}{[y^2+(t-x)^2][y^2+(t+x)^2]}\\&=\mathbf{O}\bigg(\frac{1}{y^2+(t+x)^2}\bigg)\end{aligned}$$

and then

$$\begin{aligned}&\int_\delta^\infty \frac{n_S(t)}{t}\left[\frac{t-x}{y^2+(t-x)^2}-\frac{t+x}{y^2+(t+x)^2}\right]\mathrm{d}t\\&\quad=\mathbf{O}\left(\int_\delta^\infty\frac{\mathrm{d}t}{y^2+(t+x)^2}\right)=\mathbf{O}(1).\end{aligned}$$

□

Finally we need a classical estimate of Levinson and a Phragmén–Lindelöf-type result:

Lemma 2.8. (Levinson [8]) *Let $z_n,\ n\geqslant 1$ be complex numbers with*

$$\lim_{n\to\infty}\frac{n}{z_n}=D\in\mathbb{R} \tag{2.19}$$

Suppose further that for some $c>0$

$$|z_n-z_m|\geqslant c|n-m|. \tag{2.20}$$

Let

$$F(z)=\prod_1^\infty\Big(1-\frac{z^2}{z_n^2}\Big);$$

then for any $\varepsilon>0$, as $r=|z|\to\infty$

$$F\big(re^{i\varphi}\big) \;=\; \mathbf{O}\big(e^{\pi Dr|\sin\varphi|+\varepsilon r}\big) \tag{2.21}$$

and

$$\frac{1}{F(re^{i\varphi})} \;=\; \mathbf{O}\left(e^{-\pi Dr|\sin\varphi|+\varepsilon r}\right)\ \text{if}\ \big|re^{i\varphi}-z_n\big|\geqslant\frac{1}{8}\ c. \tag{2.22}$$

Lemma 2.9. (Levin [11]) *Let $F(z)$ be an entire function of zero exponential type i.e.*

$$\limsup_{r\to\infty}\frac{\ln M(r)}{r}\leqslant 0,\quad M(r)=\max_\varphi\big|F(re^{i\varphi})\big|. \tag{2.23}$$

If $F(z)$ is bounded along a line, then $F(z)$ is constant. In particular, if $F(z)\to 0$ when $|z|\to\infty$ along a line then $F(z)\equiv 0$.

Proof of Theorem 1.9.
Step 1. Introduce the function

$$F(z)=\frac{v_1(a,z)v_2^*(a,z)-v_1^*(a,z)v_2(a,z)}{\prod_{j=1}^N p_j(z)}$$

where

$$p_j(z) = \text{p.v.} \prod_{n\in S_j} \Big(1 - \frac{z}{\lambda_n^{(j)}}\Big).$$

If $\lambda_n^{(j)} = 0$ we write z instead of $1 - \frac{z}{\lambda_n^{(j)}}$. The denominator of $F(z)$ has only simple zeros since the eigenvalues $\lambda_n^{(j)}$ are generated with a fixed right boundary condition. The values $\lambda_n^{(j)}, n \in S_j$ are zeros of the numerator, too, so $F(z)$ is an entire function. Indeed, denote by $(u_{1,j}(x,\lambda), u_{2,j}(x,\lambda))$ the solution of (2.1) with

$$u_{1,j}(0,\lambda) = \sin\alpha_j, \quad u_{2,j}(0,\lambda) = -\cos\alpha_j.$$

Since $V^* = V$ on $[0,a]$,

$$u_{1,j}^*(a,\lambda) = u_{1,j}(a,\lambda), \quad u_{2,j}^*(a,\lambda) = u_{2,j}(a,\lambda).$$

Since $\lambda_n^{(j)} = \lambda_n^{(j)^*}$ is an eigenvalue, the vectors

$$\Big(u_{1,j}(x,\lambda_n^{(j)}),\ u_{2,j}(x,\lambda_n^{(j)})\Big) \qquad \text{and} \qquad \Big(v_1(x,\lambda_n^{(j)}),\ v_2(x,\lambda_n^{(j)})\Big),$$

on the one hand, and

$$\Big(u_{1,j}^*(x,\lambda_n^{(j)}),\ u_{2,j}^*(x,\lambda_n^{(j)})\Big) \qquad \text{and} \qquad \Big(v_1^*(x,\lambda_n^{(j)}),\ v_2^*(x,\lambda_n^{(j)})\Big),$$

on the other hand are parallel. Hence

$$\frac{v_2^*\left(a,\lambda_n^{(j)}\right)}{v_1^*\left(a,\lambda_n^{(j)}\right)} = \frac{u_{2,j}^*\left(a,\lambda_n^{(j)}\right)}{u_{1,j}^*\left(a,\lambda_n^{(j)}\right)} = \frac{u_{2,j}\left(a,\lambda_n^{(j)}\right)}{u_{1,j}\left(a,\lambda_n^{(j)}\right)} = \frac{v_2\left(a,\lambda_n^{(j)}\right)}{v_1\left(a,\lambda_n^{(j)}\right)}.$$

So $F(z)$ is an entire function indeed.
Step 2. We estimate the numerator of $F(z)$ using (2.3), (2.4):

$$\begin{aligned}
&v_1(a,z)v_2^*(a,z) - v_1^*(a,z)v_2(a,z)\\
=&\sin\Big(z(\pi-a) - \int_a^\pi V - \beta\Big)\cos\Big(z(\pi-a) - \int_a^\pi V^* - \beta\Big)\\
&-\sin\Big(z(\pi-a) - \int_a^\pi V^* - \beta\Big)\cos\Big(z(\pi-a) - \int_a^\pi V - \beta\Big)\\
&+\mathbf{O}\Big(\frac{e^{2|\Im z|(\pi-a)}}{|z|}\Big)\\
=&\sin\int_a^\pi (V^* - V) + \mathbf{O}\left(\frac{e^{2|\Im z|(\pi-a)}}{|z|}\right).
\end{aligned}$$

From $a < \pi$ we know that this function has infinitely many real zeros and the zeros are not bounded. This is compatible with the above estimate only when

$$\sin\int_a^\pi (V^* - V) = 0$$

and then

$$v_1(a,z)v_2^*(a,z) - v_1^*(a,z)v_2(a,z) = \mathbf{O}\left(\frac{e^{2|\Im z|(\pi-a)}}{|z|}\right).$$

Step 3. We estimate the denominator of $F(z)$. By Lemma 2.6

$$|p_j(z)| \asymp |\widehat{p}_j(z)| \text{ if } \left|z_n^{(j)}\right| \geqslant \delta, \ \left|z - n - \frac{\vartheta_j}{\pi}\right| \geqslant \delta \text{ for } n \in S_j,$$

where

$$\widehat{p}_j(z) = \text{p.v.} \prod_{n\in S_j} \left(1 - \frac{z}{n + \frac{\vartheta_j}{\pi}}\right), \quad \vartheta_j = \beta - \alpha_j + \int_0^\pi V.$$

Again, if $n + \frac{\vartheta_j}{\pi} = 0$, we substitute $1 - \frac{z}{n+\frac{\vartheta_j}{\pi}}$ by z. In calculating $\widehat{p}_j\left(z + \frac{\vartheta_j}{\pi}\right)$ we use

$$1 - \frac{z + \frac{\vartheta_j}{\pi}}{n + \frac{\vartheta_j}{\pi}} = \frac{n - z}{n + \frac{\vartheta_j}{\pi}} = \left(1 - \frac{z}{n}\right) \frac{n}{n + \frac{\vartheta_j}{\pi}} = \left(1 - \frac{z}{n}\right)\left(1 - \frac{\frac{\vartheta_j}{\pi}}{n + \frac{\vartheta_j}{\pi}}\right).$$

to obtain

$$\widehat{p}_j\left(z + \frac{\vartheta_j}{\pi}\right) = \widetilde{p}_j(z) \cdot c_j, \quad \widetilde{p}_j(z) = \text{p.v.} \prod_{n\in S_j} \left(1 - \frac{z}{n}\right).$$

We handle the case $n = 0 \in S_j$ as above. Arrange the values $n \in S_j$ in an increasing sequence z_k. Since $n_j(z_k) = k + \mathbf{O}(1)$, we have

$$\frac{k}{z_k} = \frac{n_j(z_k)}{z_k} + \mathbf{o}(1) \longrightarrow \gamma_j \quad (k \to \infty).$$

Now the almost symmetric property of S_j implies a lower estimate by Lemma 2.8: for every $\varepsilon > 0$ there exists a $c = c(\varepsilon, \delta) > 0$ such that

$$|\widetilde{p}_j(z)| \geqslant c e^{\pi\gamma_j|\Im z| - \varepsilon|z|} \quad \text{if} \quad |z - n| \geqslant \delta \quad \text{for} \quad n \in S_j.$$

(If the product $\widetilde{p}_j$ is finite, this estimate follows immediately). By the above considerations

$$|p_j(z)| \asymp |\widehat{p}_j(z)| \asymp \left|\widetilde{p}_j\left(z - \frac{\vartheta_j}{\pi}\right)\right| \geqslant c e^{\pi\gamma_j|\Im z| - 2\varepsilon|z|}$$

for $|z|$ large enough, if $\left|z - n - \frac{\vartheta_j}{\pi}\right| \geqslant \delta$, $\left|z - \lambda_n^{(j)}\right| \geqslant \delta$ hold for $n \in S_j$. Hence the whole denominator of F has a lower estimate

$$\prod_{j=1}^N |p_j(z)| \geqslant c e^{\pi \sum\limits_j \gamma_j|\Im z| - 2\varepsilon N|z|} \geqslant c e^{2(\pi - a)|\Im z| - 2\varepsilon N|z|}.$$

By Step 2 this means that

$$|F(z)| \leqslant \ c e^{(2N+1)\varepsilon|z|} \text{ for } |z| \text{ large enough, if } \left|z - n - \frac{\vartheta_j}{\pi}\right| \geqslant \delta,$$
$$\left|z - \lambda_n^{(j)}\right| \geqslant \delta \quad \text{hold for} \quad n \in S_j.$$

Here $\varepsilon > 0$ is arbitrary and $\delta > 0$ can be chosen so small that the excluded circles of radius δ are disjoint for large $|n|$. Consequently the maximum modulus principle shows that

$$|F(z)| \leqslant c e^{(2N+1)\varepsilon|z|} \quad z \in \mathbb{C}.$$

This means that $F(z)$ is of zero exponential type.

Step 4. We show that

$$F(iy) \to 0 \quad \text{for} \quad |y| \to \infty.$$

In Step 3 we proved that

$$|p_j(iy)| \asymp |\widehat{p}_j(iy)| \asymp \left|\widetilde{p}_j\left(iy - \frac{\vartheta_j}{\pi}\right)\right|, \qquad |y| \to \infty$$

Furthermore, Lemma 2.7 states

$$\left|\widetilde{p}_j\left(iy - \frac{\vartheta_j}{\pi}\right)\right| \asymp |\widetilde{p}_j(iy)|, \qquad |y| \to \infty.$$

Consequently the denominator of F can be estimated as follows

$$\prod_{j=1}^{N} |p_j(iy)| \asymp |y|^d \exp\left(\int_{-\infty}^{\infty} \frac{\sum \widetilde{n}_j(t)}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t\right)$$

if $0 \in S_j$ is fulfilled d times. Now we have

$$\begin{aligned}&\int_1^{\infty} \frac{[t]}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t + \int_{-\infty}^{-1} \frac{-[-t]}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t\\ =&\ln\left|\frac{\sin(\pi iy)}{\pi iy}\right| = \pi|y| - \ln|y| + \mathbf{O}(1);\end{aligned}$$

because of the known formula

$$\frac{\sin \pi z}{\pi z} = \prod_{n=1}^{\infty}\left(1 - \frac{z^2}{n^2}\right)$$

and (2.11). On the other hand

$$\begin{aligned}&\int_1^{\infty} \frac{1}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t = \int_1^{\infty}\left(\frac{1}{t} - \frac{t}{y^2+t^2}\right)\mathrm{d}t\\ =&\frac{1}{2}\ln(y^2+1) = \ln|y| + \mathbf{O}(1),\end{aligned}$$

and analogously

$$\int_{-\infty}^{-1} \frac{1}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t = -\ln|y| + \mathbf{O}(1).$$

From inequality (1.19) we infer

$$\begin{aligned}&\int_{-\infty}^{\infty} \frac{\sum \widetilde{n}_j(t)}{t}\ \frac{y^2}{y^1+t^2}\mathrm{d}t \geqslant 2(\pi-a)|y|\\ &+\left(-2\left(1-\frac{a}{\pi}\right) + \left(2\left(1-\frac{a}{\pi}\right) - d + \mu - 1 + \varepsilon\right) - \mu\right)\ln|y| + \mathbf{O}(1)\\ =\ &2(\pi-a)|y| + (-1+\varepsilon-d)\ln|y| + \mathbf{O}(1)\end{aligned}$$

and then

$$\prod_{j=1}^{N} \left|p_j(iy)\right| \geqslant c|y|^{-1+\varepsilon} e^{2(\pi-a)|y|} \qquad (|y| \to \infty).$$

By Step 2

$$v_1(a,iy)v_2^*(a,iy) - v_1^*(a,iy)v_2(a,iy) \leqslant c\frac{e^{2(\pi-a)|y|}}{|y|}$$

and then

$$F(iy) = \mathbf{O}\big(|y|^{-\epsilon}\big) \to 0 \quad |y| \to \infty.$$

Now Lemma 2.9 implies

$$F(z) \equiv 0$$

or

$$\frac{v_2(a,z)}{v_1(a,z)} \equiv \frac{v_2^*(a,z)}{v_1^*(a,z)}.$$

From $V^* = V$ on $[0,a]$ it follows that

$$m(z) = \frac{v_2(0,z)}{v_1(0,z)} = \frac{v_2^*(0,z)}{v_1^*(0,z)} = m^*(z).$$

By Theorem 1.3 this implies $V^* = V$ on $[0,\pi]$. □

Proof of Theorem 1.8

It is analogous to the above proof, so we only mention the main points. Since infinitely many eigenvalues of q and of q^* are the same, from a well-known eigenfunction asymptotics (see e.g. [10], Ch.I.) we get that

$$\int_0^\pi (q^* - q) = 0. \tag{2.24}$$

Again by the standard eigenfunction asymptotics method of [10] we get

$$\begin{aligned} v(x,\lambda) &= \sin\beta\cos s(\pi - x) + \left(\cos\beta + \sin\beta \cdot \int_x^\pi \frac{q}{2}\right) \frac{\sin s(\pi - x)}{s} \\ &\quad + \mathbf{o}\left(\frac{e^{|\Im s|(\pi - x)}}{|s|}\right), \\ v'(x,\lambda) &= \sin\beta \cdot s \cdot \sin s(\pi - x) - \left(\cos\beta + \sin\beta \cdot \int_x^\pi \frac{q}{2}\right) \cos s(\pi - x) \\ &\quad + \mathbf{o}\left(e^{|\Im s|(\pi - x)}\right), \end{aligned}$$

where $\lambda = s^2$ and $s \to \infty$ along any ray $\arg s = \text{const}$. Analogously, for $\sin\beta = 0$

$$\begin{aligned} v(x,\lambda) &= \cos\beta \left(\frac{\sin s(\pi - x)}{s} - \frac{\cos s(\pi - x)}{s^2} \int_x^\pi \frac{q}{2}\right) \\ &\quad + \mathbf{o}\left(\frac{e^{|\Im s|(\pi - x)}}{|s|^2}\right), \\ v'(x,\lambda) &= -\cos\beta \left(\cos s(\pi - x) + \frac{\sin s(\pi - x)}{s} \int_x^\pi \frac{q}{2}\right) \\ &\quad + \mathbf{o}\left(\frac{e^{|\Im s|(\pi - x)}}{|s|}\right). \end{aligned}$$

Combining this with (2.24), we finally obtain

$$\begin{aligned} &v(a,z)v^{*'}(a,z) - v^*(a,z)v'(a,z) \\ &= \begin{cases} \mathbf{o}\left(e^{2|\Im\sqrt{z}|(\pi - a)}\right) & \text{if} \quad \sin\beta \neq 0 \\ \mathbf{o}\left(\frac{e^{2|\Im\sqrt{z}|(\pi - a)}}{|z|}\right) & \text{if} \quad \sin\beta = 0. \end{cases} \end{aligned} \tag{2.25}$$

The denominator of $F(z)$ can be estimated by

$$\ln|\prod_{j=1}^{N} p_j(iy)| = \int_1^\infty \frac{\sum_{j=1}^N n_j(\sqrt{t})}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t + \mathbf{O}(1).$$

Now

$$\int_1^\infty \frac{[\sqrt{t}]}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t = \ln\Big|\prod_{n=1}^{\infty}\Big(1-\frac{iy}{n^2}\Big)\Big| + \mathbf{O}(1)$$

$$= \ln\Big|\frac{\sin\pi\sqrt{iy}}{\pi\sqrt{iy}}\Big| + \mathbf{O}(1) = \frac{\pi}{\sqrt{2}}\sqrt{|y|} - \frac{1}{2}\ln|y| + \mathbf{O}(1),$$

$$\int_1^\infty \frac{[\sqrt{t}+\frac{1}{2}]}{t}\ \frac{y^2}{y^2+t^2}\mathrm{d}t = \ln\Big|\prod_{n=0}^{\infty}\Big(1-\frac{iy}{\big(n+\frac{1}{2}\big)^2}\Big)\Big| + \mathbf{O}(1)$$

$$= \ln|\cos\pi\sqrt{iy}| + \mathbf{O}(1) = \frac{\pi}{\sqrt{2}}\sqrt{|y|} + \mathbf{O}(1).$$

So by (1.17) in the case $\sin\beta \neq 0$ we obtain that

$$\ln\Big|\prod_{j=1}^{N} p_j(iy)\Big| \geqslant 2(\pi-a)\sqrt{\frac{|y|}{2}} + \mathbf{O}(1)$$

which means by (2.25) that

$$F(iy) \to 0 \qquad (y\to\infty). \tag{2.26}$$

If $\sin\beta = 0$, then similarly

$$\ln\Big|\prod_{j-1}^{N} p_j(iy)\Big| \geqslant 2(\pi-a)\sqrt{\frac{|y|}{2}} - \ln|y| + \mathbf{O}(1)$$

and from (2.25) we get (2.26). Since (2.26) implies $F(z)\equiv 0$, we are ready. □

References

1. G. Borg, *Eine Umkehrung der Sturm-Liouvilleschen Eigenwertaufgabe*, Acta Math. **78** (1946), 1–96.
2. G. Borg, *Uniqueness theorems in the spectral theory of* $y^{"} + (\lambda - q(x))y = 0$, Proc. 11th Scandinavian Congress of Mathematicians, Johan Grundt Tanums Forlag, Oslo, 1952, pp. 276–287.
3. F. Gesztesy, R. del Rio and B. Simon, *Inverse spectral analysis with partial information on the potential, III. Updating boundary conditions*, Intl. Math. Research Notices **15** (1997), 751–758.
4. F. Gesztesy, R. del Rio and B. Simon, *Corrections and Addendum to "Inverse spectral analysis with partial information on the potential, III. Updating boundary conditions"*, Intl. Math. Research Notices **11** (1999), 623–625.
5. F. Gesztesy and B. Simon, *Inverse spectral analysis with partial information on the potential, II. The case of discrete spectrum*, Trans. Amer. Math. Soc. **352** (2000), 2765–2787.
6. F. Gesztesy and B. Simon, *On the determination of the potential from three spectra*, Trans. Amer. Math. Soc. **189(2)** (1999), 85–92.
7. H. Hochstadt and B. Lieberman, *An inverse Sturm-Liouville problem with mixed given data*, SIAM J. Appl. Math. **34** (1978), 676–680.
8. N. Levinson, *Gap and density theorems*, AMS Coll. Publ., 1940, New York.
9. B. M. Levitan and I. S. Sargsjan, *Sturm-Liouville and Dirac operators (in Russian)*, Nauka, Moscow 1988.
10. B. M. Levitan and I. S. Sargsjan, *Introduction to spectral theory (in Russian)*, Nauka, Moscow 1970.

11. B. Ja. Levin, *Distribution of zeros of entire functions (in Russian)*, Gostekhizdat, Moscow 1956.
12. V.A. Marchenko, *Certain problems in the theory of second order differential operators (in Russian)*, Dokl. Akad. Nauk. SSSR, **72** (1950), 457–460.
13. B.A. Watson, *Inverse spectral problems for weighted Dirac systems*, Inverse Problems, **15(3)** (1999), 793–805.

Budapest University of
Technology and Economics
Institute of Mathematics
H 1111 Budapest, Műegyetem rkp. 3-9.
E-mail address: horvath@math.bme.hu