

Probability 1
CEU Budapest, fall semester 2016
Imre Péter Tóth

Homework sheet 8 – due on 06.12.2013 – and exercises for practice

- 8.1 (**homework**) As in Exercise 6.5, let S_n be a simple asymmetric random walk starting from $S_0 = 0$, and let τ be the hitting time for the set $H \subset \mathbb{N}$. You have seen that $M_n := \left(\frac{q}{p}\right)^{S_n}$ is a martingale and τ is a stopping time.
- Now assume that $p > \frac{1}{2}$ and $H = \{a, b\}$ where $a, b \in \mathbb{N}$. Let $p_a = \mathbb{P}(S_\tau = a)$ and $p_b = \mathbb{P}(S_\tau = b)$. What does the optional stopping theorem say about p_a and p_b ,
- (a) when $a = -5$ and $b = 7$?
 - (b) when $a = 5$ and $b = 7$?
- 8.2 Let $a, b \in \mathbb{N}$ with $a < 0 < b$. Let S_n be a simple symmetric random walk with $S_0 = 0$ and let τ be the first hitting time for $\{a, b\}$. Apply the optional stopping theorem to the martingale S_n to find the hitting probabilities $p_a = \mathbb{P}(S_\tau = a)$ and $p_b = \mathbb{P}(S_\tau = b)$.
- 8.3 Let $a, b \in \mathbb{N}$ with $a < 0 < b$. Let S_n be a simple asymmetric random walk with $p := \mathbb{P}(\text{jump to the right}) \neq \frac{1}{2}$ and $S_0 = 0$. Let τ be the first hitting time for $\{a, b\}$. Apply the optional stopping theorem to the martingale $S_n - n(p - q)$ and the result of Exercise 1 to find $\mathbb{E}\tau$.
- 8.4 Let $a, b \in \mathbb{N}$ with $a < 0 < b$. Let S_n be a simple symmetric random walk with $S_0 = 0$. Let τ be the first hitting time for $\{a, b\}$. Apply the optional stopping theorem to the martingale $S_n^2 - n$ and the result of Exercise 2 to find $\mathbb{E}\tau$.
- 8.5 *Life, the Universe, and Everything.* Arthur decides to keep rolling a fair die until he manages to roll two 6-es consecutively. What is the expected number of rolls he needs?
- 8.6 (**homework**) Bob keeps tossing a fair coin and makes notes of the results: he writes “H” for heads and “T” for tails. Calculate the expected number of tosses
- a.) until the character sequence “HTHT” shows up,
 - b.) until the character sequence “THTT” shows up.
- 8.7 Alice and Bob keep tossing a fair coin until either the word $A := \text{“HTHT”}$ or the word $B := \text{“THTT”}$ shows up. If the word appearing first is A , then Alice wins, and if B , then Bob. Introduce the notation $p_A := \mathbb{P}(\text{Alice wins})$, $p_B := \mathbb{P}(\text{Bob wins})$. Let τ be the random time when the game ends.
- a.) Think of a casino, as in the solution of the ABRACADABRA problem [2], where all players bet for (consecutive letters of) the word A . Using the capital of this casino as a martingale, express $\mathbb{E}\tau$ using p_A and p_B .
 - b.) Now think of another casino, where all players bet for (consecutive letters of) the word B . Using the capital of this other casino as a martingale, get another expression for $\mathbb{E}\tau$ using p_A and p_B .
 - c.) Solve the system of equations formed by the two equations above, to calculate $\mathbb{E}\tau$, p_A and p_B .

8.8 A monkey keeps pressing keys of a typewriter with 26 keys printing the letters of the English alphabet, uniformly and independently of the past, until the word “ABRACADABRA” shows up. Denote this random time by τ . Beside the monkey – as in the original ABRACADABRA solution [2]– operates a casino where players can always bet for the next key pressed in a fair game: if their guess is wrong, they lose their bet entirely, while if it is correct, they lose it and get back 26 times more.

Before every keypress, a new player arrives, who will bet all his money first on “A”, then on “B”, then on “R”, etc. through the ABRACADABRA sequence, as long as he keeps winning or the game ends. (If he loses once, he goes home immediately.) This is just like in the original ABRACADABRA solution.

However, the later a player arrives, the less money he has to play with: there is some fixed $z \in (0, 1)$ such that the n -th player arrives with $\$z^{n-1}$.

Show that the fortune of the casino is a martingale, and use the optional stopping theorem to calculate the generating function of τ .

8.9 Let N, X_1, X_2, X_3, \dots be independent, and let them all have (optimistic) geometric distribution with parameter $p = \frac{1}{6}$. Calculate the expectation of

$$S =: \sum_{k=1}^N (X_k + 1).$$

What has this got to do with Exercise 5?

Hint: use the generating function method, or simply apply the theorem we had about sums with random number of terms.

8.10 (**homework**) Durrett [1], Exercise 8.1.3

8.11 Durrett [1], Exercise 8.2.3

8.12 (*bonus for those who are interested*) It is not hard to show that if ξ is a standard Gaussian random variable and $x \geq 1$, then

$$\mathbb{P}(|X| \geq x) \leq \sqrt{\frac{2}{\pi}} e^{-\frac{x^2}{2}}.$$

Use this to show that if ξ_1, ξ_2, \dots are i.i.d. standard Gaussian, then, with probability 1, the event $\{|\xi_n| > 2 \ln n\}$ occurs for at most finitely many n -s.

8.13 (*bonus for those who are interested*) *Paul Lévy construction of the Wiener process.* In a possible construction of the Wiener process (or Brownian motion) on $[0, 1]$ we define a sequence of piecewise linear continuous random functions so that we first define f_n at dyadic rationals that are multiples of $\frac{1}{2^n}$, inheriting every second value (at multiples of $\frac{1}{2^{n-1}}$) from f_{n-1} , and setting the values at the remaining points (of the form $\frac{2k-1}{2^n}$) to be the average of the two neighbouring values, plus an independent Gaussian random value with mean 0 and variance $\frac{1}{4^n}$. Then we extend f_n to $[0, 1]$ piecewise linearly.

Formally: we take independent standard Gaussian random variables ξ_0 and $\xi_{n,k}$ where $n = 1, 2, \dots$ and $k = 1, 2, \dots, 2^{n-1}$. Then

- In the 0th step we fix $f_0(0) = 0$ and $f_0(1) = \xi_0$. We connect these two values linearly.
- In the 1st step we leave $f_1(0) = f_0(0)$ and $f_1(1) = f_0(1)$, but also set $f_1(\frac{1}{2}) = f_0(\frac{1}{2}) + \frac{1}{2}\xi_{1,1}$. We connect these three values linearly.

- ...in the n th step we leave $f_n\left(\frac{k}{2^{n-1}}\right) = f_{n-1}\left(\frac{k}{2^{n-1}}\right)$ for $k = 0, 1, \dots, 2^{n-1}$, but also set $f_n\left(\frac{k-\frac{1}{2}}{2^{n-1}}\right) = f_{n-1}\left(\frac{k-\frac{1}{2}}{2^{n-1}}\right) + \frac{1}{2^n}\xi_{n,k}$ for $k = 1, \dots, 2^{n-1}$. We connect these $2^n + 1$ values linearly.

Notice that, in this construction, the difference $g_n := f_{n+1} - f_n$ is the sum of 2^n “tent” maps with disjoint supports and i.i.d. Gaussian “heights”.

- (a) Use the statement of Exercise 12 to show that, with probability 1, the series

$$\lim_{n \rightarrow \infty} f_n = f_0 + \sum_{n=0}^{\infty} g_n$$

is uniformly absolutely convergent.

- (b) Check that the limit is a Wiener process.

References

- [1] Durrett, R. *Probability: Theory and Examples*. **4th** edition, Cambridge University Press (2010)
- [2] Ai, Di. *Martingales and the ABRACADABRA problem*.
<http://math.uchicago.edu/~may/VIGRE/VIGRE2011/REUPapers/Ai.pdf> (2011)