Problem Set 1 Brownian Motion: Construction and Basic Properties

1.1 Let

 $\varphi : \mathbb{R} \to \mathbb{R}_+, \quad \varphi(x) := \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \quad \text{be the standard normal density function,}$ $\Phi : \mathbb{R} \to [0, 1], \quad \Phi(x) := \int_{-\infty}^x \varphi(y) dy, \quad \text{be the standard normal distribution function.}$

Prove that for any x > 0

$$\left(\frac{1}{x} - \frac{1}{x^3}\right)\varphi(x) < 1 - \Phi(x) < \frac{1}{x}\varphi(x).$$

Hint: Compare the derivatives.

1.2 Let $X_1^{(n)}, X_2^{(n)}, \ldots, X_n^{(n)}$ be i.i.d. normal random variables with

$$\mathbf{E}\left(X_{j}^{(n)}\right) = 0, \qquad \mathbf{Var}\left(X_{j}^{(n)}\right) = \frac{1}{n}, \qquad j = 1, \dots, n.$$

Define the stochastic process $t \mapsto B^{(n)}(t), t \in [0, 1]$ as follows:

$$B^{(n)}(t) := \sum_{j=1}^{\lfloor nt \rfloor} X_j^{(n)}.$$

(a) Compute the expectations and covariances

$$\mathbf{E}(B^{(n)}(t)) = ?, \quad \mathbf{Cov}(B^{(n)}(t), B^{(n)}(s)) = ?, \quad s, t \in [0, 1],$$

and their limits as $n \to \infty$.

(b) What is the joint distribution of the random variables $\{B^{(n)}(t) : t \in [0,1]\}$?

(c) Let

$$\delta_n := \max\left\{ \left| B^{(n)}(t+) - B^{(n)}(t-) \right| : t \in [0,1] \right\}.$$

(In plain words: δ_n is the largest jump discontinuity of the process $\{B^{(n)}(t) : t \in [0,1]\}$.)

Prove that for any fixed $\varepsilon > 0$,

$$\lim_{n \to \infty} \mathbf{P} \left(\delta_n \ge \varepsilon \right) = 0.$$

Hint: Note that $\delta_n = \max_{1 \le j \le n} |X_j^{(n)}|$ and use the upper bound from problem 1.1.

(d) Use your favourite program package (R, Matlab, Mathematica, Maple, ...) to simulate the process and draw (print) its trajectory for n = 100, 1000, and 10000.

1.3 Let $Y_1^{(n)}, Y_2^{(n)}, \ldots, Y_n^{(n)}$ be i.i.d. Poisson random variables with parameter 1/n. So,

$$\mathbf{E}\left(Y_{j}^{(n)}\right) = \frac{1}{n}, \qquad \mathbf{Var}\left(Y_{j}^{(n)}\right) = \frac{1}{n}, \qquad j = 1, \dots, n.$$

Define the stochastic process $t \mapsto B^{(n)}(t), t \in [0, 1]$ as follows:

$$Z^{(n)}(t) := \sum_{j=1}^{\lfloor nt \rfloor} \left(Y_j^{(n)} - \frac{1}{n} \right)$$

(a) Compute the expectations and covariances

$$\mathbf{E}(Z^{(n)}(t)) = ?, \quad \mathbf{Cov}(Z^{(n)}(t), Z^{(n)}(s)) = ?, \quad s, t \in [0, 1],$$

and their limits as $n \to \infty$.

(b) What is the joint distribution of the random variables $\{Z^{(n)}(t) : t \in [0,1]\}$? Explain in plain words.

(c) Let

$$\delta_n := \max\left\{ \left| Z^{(n)}(t+) - Z^{(n)}(t-) \right| : t \in [0,1] \right\}.$$

(In plain words: δ_n is the largest jump discontinuity of the process $\{Z^{(n)}(t) : t \in [0,1]\}$.)

Compute, for $\varepsilon > 0$ fixed,

$$\lim_{n \to \infty} \mathbf{P} \left(\delta_n \ge \varepsilon \right).$$

Hint: Note that $\delta_n = \max_{1 \le j \le n} |Y_j^{(n)}|$ and use all you know about Poisson random variables.

(d) Use your favourite program package (R, Matlab, Mathematica, Maple, ...) to simulate the process and draw (print) its trajectory for n = 100, 1000, and 10000.

- **1.4** Interpret the results of problems 1.2, respectively, 1.3.
- **1.5** (a) Let Y_1, Y_2, \ldots, Y_n be random variables with $\mathbf{E}(Y_j) = 0$ and $\mathbf{Cov}(Y_i, Y_j) =: c_{i,j}$. Assume that the covariance matrix $C := (c_{i,j})_{i,j=1}^n$ is non-degenerate, $\det(C) \neq 0$. Prove that the random variables Y_1, Y_2, \ldots, Y_n are *jointly Gaussian* if and only if there exist i.i.d. $\mathcal{N}(0, 1)$ -distributed random variables X_1, X_2, \ldots, X_n and real coefficients $(a_{i,j})_{i,j=1}^n$ such that

$$Y_i = \sum_{j=1}^n a_{ij} X_j.$$

Hint: Express the matrix $A = (a_{i,j})_{i,j=1}^n$ from the covariance matrix $C = (c_{i,j})_{i,j=1}^n$. (b) Let $t \mapsto B(t)$ be standard 1d Brownian motion and $0 \le t_1 \le t_2 \le \cdots \le t_n$. Explain why it follows from the definition of Brownian motion (i.e. independent and Gaussian increments) that the random variables $B(t_1), B(t_2), \ldots, B(t_n)$ have jointly Gaussian distribution.

- (c) Determine the covariance matrix of the random variables $B(t_1), B(t_2), \ldots, B(t_n)$.
- **1.6** Let $t \mapsto B(t)$ be standard 1*d* Brownian motion. Prove that the following processes are also standard 1*d* Brownian motions:
 - (a) The rescaled process: $X(t) := a^{-1/2}B(at)$, where a > 0 is fixed parameter.
 - (b) The time reversed process: Y(t) := tB(1/t).
 - (c) The backwards process: Z(t) := B(T) B(T t), where T > 0 is fixed and $t \in [0, T]$.

Hint: Prove that the processes X(t), Y(t), Z(t) are Gaussian and compute their covariances.

- **1.7** For j = 1, ..., n, let $t \mapsto B_j(t)$, be independent 1*d* Brownian motions with variance σ_j^2 , and a_j fixed real numbers. Prove that the process $t \mapsto Z(t) := \sum_{j=1}^n a_j B_j(t)$ is also a 1*d* Brownian motion. Determine the variance of the process Z(t).
- **1.8** Let $t \mapsto B(t)$ be standard 1*d* Brownian motion. Determine (without painful computations) the conditional probability

$$\mathbf{P}(B(2) > 0 \mid B(1) > 0).$$

- **1.9** Show that 1*d* Brownian motion changes sign infinitely many times in any time interval $[0, \delta]$ of positive length δ .
- 1.10 The Brownian meander process.

(a) Let $\varepsilon > 0$ be fixed. Using the *reflection principle* prove that for any x > 0, t > 0

$$\mathbf{P}(B(t) \ge x - \varepsilon \mid \min_{0 \le s \le t} B(s) \ge -\varepsilon) = \frac{\Phi((-x + \varepsilon)/\sqrt{t}) - \Phi((-x - \varepsilon)/\sqrt{t})}{2\Phi(\varepsilon/\sqrt{t}) - 1}.$$
 (*)

(b) Letting $\varepsilon \to 0$ in the previous formula prove that the conditional density of B(t), given $\{B(s) \ge 0 : s \in [0, t]\}$ is

$$\frac{x}{t} \exp\{-x^2/(2t)\} 1_{\{x>0\}}.$$

Remark: Note that the probability of the condition is zero (see problem 1.9). Brownian motion conditioned to stay positive is called *Brownian meander*.

1.11 On the Hilbert space $\mathcal{L}^2([0, 1], dx)$ define the self-adjoint compact (actually: Hilbert-Schmidt) operator

$$Kf(t) := \int_0^1 \min\{t, s\} f(s) ds$$

Prove that

$$\lambda_n = \frac{4}{\pi^2 (2n-1)^2}, \qquad \psi_n(t) = \sqrt{2} \sin\left(\frac{\pi (2n-1)}{2}t\right), \qquad n = 1, 2, \dots$$

are eigenvalues and eigenvectors of the operator K.

1.12 Let ξ be a standard normal random variable and define, for $\lambda < 1$

$$\psi(\lambda) := \log \mathbf{E} \left(\exp\{\lambda(\xi^2 - 1)/2\} \right).$$

Prove that

$$\psi(\lambda) = -\frac{1}{2} \left(\log(1-\lambda) + \lambda \right),$$

and investigate the analytic properties of the function $\psi(\cdot)$ (convexity, minima, asymptotes, ...). Plot the graph of the function $\lambda \mapsto \psi(\lambda)$.

1.13 Show that the function

$$\phi : \mathbb{R}_+ \times \mathbb{R} \to \mathbb{R}, \qquad \phi(t, x) := \frac{1}{\sqrt{t}} \varphi(\frac{x}{\sqrt{t}})$$

solves the heat equation

$$\partial_t \phi(t,x) = \frac{1}{2} \partial_x^2 \phi(t,x).$$