

19th and 20th lectures

L'Hospital's rule

Theorem (L'Hospital's rule).

Assume that $a \in \overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$, I is a neighbourhood of a , the functions f and g are differentiable on $I \setminus \{a\}$ and $g(x) \neq 0$, $g'(x) \neq 0$ for all $x \in I \setminus \{a\}$. Assume moreover that

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0 \quad \text{or} \quad \lim_{x \rightarrow a} |f(x)| = \lim_{x \rightarrow a} |g(x)| = \infty.$$

$$\text{If } \exists \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = b \in \overline{\mathbb{R}} \text{ then } \exists \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = b.$$

Remark. The theorem holds for right-hand and left-hand limits as well.

Proof. We prove it in the case when $a \in \mathbb{R}$ (for right-hand limit).

$$\text{Assume that } a \in \mathbb{R}, \lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} g(x) = 0 \text{ and } \exists \lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} = b \in \mathbb{R}.$$

Extend the functions f and g such that $f(a) = g(a) = 0$ and let $x \in I$, $x > a$.

Then f and g are continuous on $[a, x]$ and differentiable on (a, x) ,

so by Cauchy's mean value theorem there exists $c \in (a, x)$ such that

$$\frac{f(x)}{g(x)} = \frac{f(x) - f(a)}{g(x) - g(a)} = \frac{f'(c)}{g'(c)}.$$

Let (x_n) be a sequence such that $x_n \rightarrow a$ and choose $c_n \in (a, x_n)$ for all n .

$$\text{Then } c_n \rightarrow a \text{ and } \frac{f(x_n)}{g(x_n)} = \frac{f'(c_n)}{g'(c_n)} \text{ for all } n \in \mathbb{N}.$$

$$\text{Therefore } \lim_{n \rightarrow \infty} \frac{f(x_n)}{g(x_n)} = \lim_{n \rightarrow \infty} \frac{f'(c_n)}{g'(c_n)} = b \text{ and by the sequential criterion for the limit, } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = b.$$

Undefined forms

Remark. The theorem can be applied for limits of the following type:

$$1) \frac{0}{0}, \frac{\infty}{\infty} : \text{L'Hospital's rule can be applied directly}$$

$$2) 0 \cdot \infty : \text{we can try the following transformations: } f(x)g(x) = \frac{f(x)}{\frac{1}{g(x)}} \text{ or } f(x)g(x) = \frac{g(x)}{\frac{1}{f(x)}}$$

$$3) \infty - \infty : h(x) = \frac{1}{f(x)}, k(x) = \frac{1}{g(x)} \implies f(x) - g(x) = \frac{1}{h(x)} - \frac{1}{k(x)} = \frac{k(x) - h(x)}{h(x)k(x)} \quad \left(\begin{array}{c} 0 \\ - \\ 0 \end{array} \right)$$

$$4) 0^0, 1^\infty, \infty^0 : (f(x))^{g(x)} = e^{g(x) \cdot \ln(f(x))}, \text{ then for the undefined form } g(x) \cdot \ln(f(x)) \text{ previous methods can be applied.}$$

Exercises

Pages 171-172 of the pdf file (first 9 examples):

https://math.bme.hu/~tasnadi/merninf_anal_1/anal1_elm.pdf

Pages 72-73 of the pdf file, exercise 26:

https://math.bme.hu/~tasnadi/merninf_anal_1/anal1_gyak.pdf

In exercises 26. g), h) the L'Hospital's rule cannot be applied.

Local properties and the derivative

Definition. Assume that $x_0 \in D_f$ and there exists $\delta > 0$ such that for all $x, y \in D_f$, if $x_0 - \delta < x < x_0 < y < x_0 + \delta$,

then $\begin{cases} f(x) \leq f(x_0) \leq f(y) \\ f(x) \geq f(x_0) \geq f(y) \\ f(x) < f(x_0) < f(y) \\ f(x) > f(x_0) > f(y) \end{cases}$. Then we say that f is $\begin{cases} \text{locally increasing} \\ \text{locally decreasing} \\ \text{strictly locally increasing} \\ \text{strictly locally decreasing} \end{cases}$ at x_0 .


Remarks. (1) If f is monotonically increasing on (a, b) , then f is locally increasing for all $x_0 \in (a, b)$.

(2) If f is locally increasing **for all** $x_0 \in (a, b)$, then f is monotonically increasing on (a, b) .


(3) However, if f is locally increasing at x_0 then it doesn't imply that there exists a neighbourhood $B(x_0, r)$ where f is monotonically increasing.

The following functions are locally increasing at $x_0 = 0$ but on any interval that contains 0, the functions are not monotonically increasing.


$$1. f(x) = \begin{cases} x \sin^2 \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$



$$2. f(x) = \begin{cases} \frac{1}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$



$$3. f(x) = \begin{cases} x & \text{if } x \in \mathbb{Q} \\ 2x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q} \end{cases}$$



Theorem. Assume that f is differentiable at x_0 .

- (1) If f is locally increasing at x_0 then $f'(x_0) \geq 0$.
- (2) If f is locally decreasing at x_0 then $f'(x_0) \leq 0$.
- (3) If $f'(x_0) > 0$ then f is strictly locally increasing at x_0 .
- (4) If $f'(x_0) < 0$ then f is strictly locally decreasing at x_0 .

Proof. (1) If f is locally increasing at x_0 then $\exists \delta > 0$ such that

$$0 < |x - x_0| < \delta \implies \frac{f(x) - f(x_0)}{x - x_0} \geq 0.$$

(If $x < x_0$ then $x - x_0 < 0$ and $f(x) - f(x_0) \leq 0$ and

if $x > x_0$ then $x - x_0 > 0$ and $f(x) - f(x_0) \geq 0$.)

$$\text{Since } f \text{ is differentiable at } x_0 \text{ then } f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \geq 0.$$

(2) Similar to case (1).

(3) If $f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} > 0$, then there exists $\delta > 0$ such that

$$\text{if } 0 < |x - x_0| < \delta \text{ then } \frac{f(x) - f(x_0)}{x - x_0} > 0.$$

$$\implies \text{if } \begin{cases} x_0 < x < x_0 + \delta \\ x_0 - \delta < x < x_0 \end{cases} \text{ then } \begin{cases} f(x) > f(x_0) \\ f(x) < f(x_0) \end{cases}$$

$\implies f$ is strictly locally increasing at x_0 .

(3) Similar to case (4).

Remarks. Assume that f is differentiable at x_0 .


(1) If f is strictly locally increasing at x_0 then it doesn't imply that $f'(x_0) > 0$.

If f is strictly locally increasing at x_0 then $f'(x_0) \geq 0$, since $\exists \delta > 0$ such that


$$0 < |x - x_0| < \delta \implies \frac{f(x) - f(x_0)}{x - x_0} > 0, \text{ but for the limit } \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \geq 0.$$

For example $f(x) = x^3$ is strictly locally increasing at $x_0 = 0$, but $f'(0) = 3x^2|_{x=0} = 0$.


1. $f(x) = x^3$



2. $f(x) = -x^3$



3. $f(x) = \begin{cases} x + x^2 \sin\left(\frac{10}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$



(2) If $f'(x_0) \geq 0$ then it doesn't imply that f is locally increasing at x_0 .

For example $f(x) = -x^3$ is not locally increasing at $x_0 = 0$, but $f'(0) = 0 \geq 0$.

(3) If $f'(x_0) > 0$ then it doesn't imply that f is monotonically increasing on an interval containing x_0 .

For example, let f be a function such that $x - x^2 \leq f(x) \leq x + x^2 \quad \forall x \implies f(0) = 0$.

$$\text{If } x > 0 \text{ then } 1 - x \leq \frac{f(x) - f(0)}{x - 0} \leq 1 + x,$$

$$\text{If } x < 0 \text{ then } 1 - x \geq \frac{f(x) - f(0)}{x - 0} \geq 1 + x, \text{ so by the sandwich theorem}$$

$$f'(0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = 1 > 0. \text{ For example, let } f(x) = \begin{cases} x + x^2 \sin\left(\frac{10}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

Darboux's theorem

Theorem. Assume that $f : [a, b] \rightarrow \mathbb{R}$ is differentiable and $f'(a) < y < f'(b)$ or $f'(b) < y < f'(a)$. Then there exists $c \in (a, b)$ such that $f'(c) = y$.

Remark. We say that f' has the intermediate value property of Darboux property.

Proof. 1) Let $g : [a, b] \rightarrow \mathbb{R}$, $g(x) = f(x) - y \cdot x \implies g$ is differentiable and $g'(x) = f'(x) - y$.

2) Assume that $f'(a) < y < f'(b) \implies g'(a) = f'(a) - y < 0 < f'(b) - y < g'(b)$

3) g is differentiable, so it is continuous on $[a, b]$

\implies by Weierstrass extreme value theorem it has a minimum and a maximum on $[a, b]$.

4) Since $\begin{cases} g'(a) < 0 \\ g'(b) > 0 \end{cases}$ then $\begin{cases} g \text{ is strictly locally decreasing at } a \\ g \text{ is strictly locally increasing at } b \end{cases}$

$\implies g$ does not have a minimum at a and b but on the open interval (a, b)

\implies there exists $c \in (a, b)$ such that g has a local minimum at c

$\implies g'(c) = 0 = f'(c) - y \implies f'(c) = y$ for some $c \in (a, b)$.

Example. The sign function or signum function is defined as $\text{sgn } x = \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0. \\ 1 & \text{if } x > 0 \end{cases}$

This function is not continuous at $x_0 = 0$, so there is no function $f : \mathbb{R} \rightarrow \mathbb{R}$ for which $f'(x) = \text{sgn } x$ on \mathbb{R} (or on any interval that contains $x_0 = 0$).

Remark. From Darboux's theorem it follows that if f' is not continuous at a point then f' cannot have a discontinuity of the first type at that point, so at least one of the one-sided limits doesn't exist or exists but is not finite $\implies f'$ has an essential discontinuity at the given point.

Statement. If f is differentiable on $[a, a + \delta)$ ($\delta > 0$) and f' has a discontinuity at a then the limit $\lim_{x \rightarrow a+0} f(x)$ doesn't exist or $\exists \lim_{x \rightarrow a+0} f(x) \notin \mathbb{R}$.

Continuously differentiable functions


Definition. Assume that I is a neighbourhood of $a \in D_f$ and f is differentiable on $I \cap D_f$.

Then f is **continuous differentiable at a** if f' is continuous at a .

f is **continuously differentiable** on A if f is continuous differentiable $\forall x \in A$.

Notation: $C^1(A) = \{f : f \text{ is continuously differentiable on } A\}$.

Example: The function $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$ is differentiable but f' is not continuous at $x_0 = 0$, since $f'(x) = \begin{cases} 2x \sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$.



Higher order derivatives

Definition. If f' is differentiable at x then we say that f is twice differentiable at x and the second derivative or second order derivative of f at x_0 is $f''(x) = (f')'(x)$.

Differentiating f repeatedly, we get the third, ..., n th derivative of f .

$$\text{Notation: } f''(x) = f^{(2)}(x) = \frac{d^2 f(x)}{dx^2}$$

$$f'''(x) = f^{(3)}(x) = \frac{d^3 f(x)}{dx^3}$$

...

$$f^{(n)}(x) = \frac{d^n f(x)}{dx^n}$$

By definition: $f^{(0)}(x) = f(x)$

Example: $f(x) = \sin x \implies f'(x) = \cos x, f''(x) = -\sin x, f'''(x) = -\cos x, f^{(4)}(x) = \sin x, \dots$
 $f(x) = e^x \implies f^{(n)}(x) = e^x \quad \forall n \in \mathbb{N}$

Investigation of differentiable functions

Monotonicity on an interval

Theorem. Assume that $f : (a, b) \rightarrow \mathbb{R}$ is differentiable. Then

- (1) f is monotonically increasing $\iff f'(x) \geq 0$ for all $x \in (a, b)$
- (2) f is monotonically decreasing $\iff f'(x) \leq 0$ for all $x \in (a, b)$
- (3) f is constant $\iff f'(x) = 0$ for all $x \in (a, b)$
- (4) $f'(x) > 0$ for all $x \in (a, b) \implies f$ is strictly monotonically increasing
- (5) $f'(x) < 0$ for all $x \in (a, b) \implies f$ is strictly monotonically decreasing

Proof. (1)

- (i) If f is monotonically increasing then f is locally monotonically increasing for all $x \in (a, b)$
 $\implies f'(x) \geq 0 \quad \forall x \in (a, b)$.
- (ii) Assume that $f'(x) \geq 0$ for all $x \in (a, b)$. Let $a < x_1 < x_2 < b$ and apply Lagrange's mean value theorem for $[x_1, x_2]$. Then there exists $c \in (x_1, x_2) \subset (a, b)$ such that

$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = f'(c) \geq 0 \implies f(x_2) \geq f(x_1)$$

Therefore if $x_1 < x_2$ then $f(x_1) \leq f(x_2)$, so f is monotonically increasing on (a, b) .

(2) Similar to case (1).

(3) f is constant $\iff f$ is monotonically increasing and decreasing
 $\iff f'(x) \geq 0$ and $f'(x) \leq 0 \quad \forall x \in (a, b) \iff f'(x) = 0 \quad \forall x \in (a, b)$

(4) and (5): similar to case (1) (ii)

Remark. Statements (4) and (5) cannot be reversed.

For example, $f(x) = x^3$ is strictly monotonically increasing on \mathbb{R} , however $f'(x) > 0$ does not hold for all $x \in \mathbb{R}$, since $f'(x) = 3x^2 \implies f'(0) = 0$.

Remark. If the domain of f is not an interval then the above theorem is not true, as the following examples show.

1) Let $f : \mathbb{R} \setminus \mathbb{Z} \rightarrow \mathbb{R}$, $f(x) = \{x\} = x - [x]$. Then f is differentiable on $\mathbb{R} \setminus \mathbb{Z}$ and $f'(x) = 1 > 0$ for all $x \in \mathbb{R} \setminus \mathbb{Z}$ but f is not monotonically increasing.

2) Let $f : \mathbb{R} \setminus \mathbb{Z} \rightarrow \mathbb{R}$, $f(x) = [x]$. Then f is differentiable on $\mathbb{R} \setminus \mathbb{Z}$ and $f'(x) = 0$ for all $x \in \mathbb{R} \setminus \mathbb{Z}$ but f is not constant.

Local extremum, sufficient conditions

Definition. If f is differentiable at x_0 and $f'(x_0) = 0$ then x_0 is a **stationary point** of f .
If $f'(x_0) = 0$ or f is not differentiable at x_0 then x_0 is a **critical point** of f .

Remark. Recall that if f is differentiable at $x_0 \in \text{int } D_f$ and f has a local extremum at x_0 then $f'(x_0) = 0$.
This is a necessary condition for the existence of a local extremum.
The next two theorems formulate sufficient conditions.

Theorem (Sufficient condition for a local extremum, first derivative test).

Assume that f is differentiable at $x_0 \in \text{int } D_f$.

If $f'(x_0) = 0$ and f' changes sign at x_0 , then f has a local extremum at x_0 .

Namely, if $f'(x_0) = 0$ and f' is (strictly) locally $\begin{cases} \text{increasing} \\ \text{decreasing} \end{cases}$ at x_0

then f has a (strict) local $\begin{cases} \text{minimum} \\ \text{maximum} \end{cases}$ at x_0 .

Proof. Assume that $f'(x_0) = 0$ and f' is locally increasing at x_0
(that is, f' changes sign from negative to positive)

$$\Rightarrow \exists \delta > 0 \text{ such that } \begin{cases} f'(x) \leq 0 & \text{if } x_0 - \delta < x < x_0 \\ f'(x) \geq 0 & \text{if } x_0 < x < x_0 + \delta \end{cases}$$

$$\Rightarrow \begin{cases} f \text{ is monotonically decreasing on } (x_0 - \delta, x_0) \\ f \text{ is monotonically increasing on } (x_0, x_0 + \delta) \end{cases}$$

$$\Rightarrow \begin{cases} f(x) \geq f(x_0) & \text{if } x_0 - \delta < x < x_0 \\ f(x) \geq f(x_0) & \text{if } x_0 < x < x_0 + \delta \end{cases} \Rightarrow f \text{ has a local minimum at } x_0.$$

Theorem (Sufficient condition for a local extremum, second derivative test).

Assume that f is twice differentiable at $x_0 \in \text{int } D_f$.

If $f'(x_0) = 0$ and $f''(x_0) \neq 0$ then f has a local extremum at x_0 .

If $\begin{cases} f''(x_0) > 0 \\ f''(x_0) < 0 \end{cases}$ then f has a strict local $\begin{cases} \text{minimum} \\ \text{maximum} \end{cases}$ at x_0 .

Proof. $f''(x_0) > 0 \Rightarrow f'$ is locally increasing at x_0 and $f'(x_0) = 0$
 \Rightarrow by the previous theorem f has a local minimum at x_0 .

Remark. The sign change of f' at x_0 is only a sufficient but not a necessary condition for the existence of a local extremum at x_0 .

$$\text{For example, if } f(x) = \begin{cases} x^2 \left(2 + \sin\left(\frac{1}{x}\right) \right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

then f is differentiable for all $x \in \mathbb{R}$. At $x = 0$:

$$f'(0) = \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{x^2 \left(2 + \sin\left(\frac{1}{x}\right) \right)}{x} = \lim_{x \rightarrow 0} x \left(2 + \sin\left(\frac{1}{x}\right) \right) = 0 \quad (\text{since it is } 0 \cdot \text{bounded}),$$

so the necessary condition holds at $x_0 = 0$.


However, in any neighbourhood of $x_0 = 0$:

f has strictly monotonic increasing and decreasing sections \implies

f' has both positive and negative values \implies

f' doesn't change sign at $x_0 = 0$.

Yet f has a local extreme value at $x_0 = 0$, and it is even an absolute minimum here.



Local extremum and higher order derivatives

Remark. If $f'(x_0) = 0$ and $f''(x_0) = 0$ then it cannot be decided whether f has a local extremum at x_0 . For example:

- 1) $f(x) = x^3$ does not have a local extremum at $x_0 = 0$,
- 2) $f(x) = x^4$ has a local minimum at $x_0 = 0$,
- 3) $f(x) = -x^4$ has a local maximum at $x_0 = 0$, and in each case $f'(0) = f''(0) = 0$.

Theorem. (1) Assume that f is $2k$ times differentiable at x_0 , $k \geq 1$.

$$\text{If } f'(x_0) = \dots = f^{(2k-1)}(x_0) = 0 \text{ and } \begin{cases} f^{(2k)}(x_0) > 0 \\ f^{(2k)}(x_0) < 0 \end{cases}$$

then f has a strict local $\begin{cases} \text{minimum} \\ \text{maximum} \end{cases}$ at x_0 .

(2) Assume that f is $2k + 1$ times differentiable at x_0 , $k \geq 1$.

If $f'(x_0) = \dots = f^{(2k)}(x_0) = 0$ and $f^{(2k+1)}(x_0) \neq 0$, then f is strictly monotonic in a neighbourhood of x_0 , so f doesn't have a local extremum at x_0 .

Remark. Part (1) in other words: If the first non-zero derivative (after the first one) has an even order then f has a local extremum at x_0 .

Proof. (1) We prove the statement for a strict local minimum by induction.

(i) If $k = 1$ then the statement is true.

(ii) Assume that the statement holds for $k - 1$ and let $g = f''$.

$$(\implies g' = f''', \dots, g^{(2k-3)} = f^{(2k-1)}, g^{(2k-2)} = f^{(2k)}.)$$

From the induction hypothesis it follows that

if $g'(x_0) = \dots = g^{(2k-3)}(x_0) = 0$ and $g^{(2k-2)}(x_0) > 0$ then the function $g = f''$ has a strict local minimum at x_0 .

(iii) We want to prove that if

$f'(x_0) = f''(x_0) = f'''(x_0) = \dots = f^{(2k-1)}(x_0) = 0$ and $f^{(2k)}(x_0) > 0$ then f has a strict local minimum at x_0 .

Since $f''(x_0) = 0$ and f'' has a strict local minimum at x_0 ,

then $\exists \delta > 0$ such that $f''(x) > 0, \forall x \in (x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$

$\implies f'$ is strictly monotonically increasing on $(x_0 - \delta, x_0 + \delta)$

$\implies f'$ is strictly locally increasing at x_0

$\implies f$ has a strict local minimum at x_0 .

(2) Assume that $f'(x_0) = f''(x_0) = \dots = f^{(2k)}(x_0) = 0$ and $f^{(2k+1)}(x_0) \neq 0$.

Let $g = f'$, then $g'(x_0) = \dots = g^{(2k-1)}(x_0) = 0$ and $g^{(2k)}(x_0) \neq 0$.

\implies by part (1), $g = f'$ has a strict local extremum at x_0 .

Since $f'(x_0) = 0$, then either $f'(x) > 0$ or $f'(x) < 0, \forall x \in (x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$

$\implies f$ is strictly monotonic on $(x_0 - \delta, x_0 + \delta)$

$\implies f$ doesn't have a local extremum at x_0 .


Example. $f(x) = x^n$ is n times differentiable,

$$f^{(k)}(x) = n(n-1)(n-2)\dots(n-k+1)x^{n-k}, \quad k = 1, 2, \dots, n-1$$

$$f^{(n)}(x) = n!$$

\implies if $x_0 = 0$, then $f'(0) = f''(0) = \dots = f^{(n-1)}(0) = 0, f^{(n)}(0) = n! > 0$

\implies at $x_0 = 0$ f has a local minimum if n is even and f doesn't have a local extremum if n is odd.



Convexity / concavity on an interval


Theorem (Necessary and sufficient condition for convexity).

If f is differentiable on the interval I , then the following statements are equivalent.

- (1) f is convex on I
- (2) $f(x) \geq f(a) + f'(a)(x - a)$ if $x, a \in I$
- (3) f' is monotonically increasing on I

Remark. The geometrical meaning of (2) is that for all $a \in I$, the graph of f lies above the tangent line at a .

Proof of (1) \Rightarrow (2):



If $a < x$ and $y \in (a, x)$ then $\exists \lambda \in (0, 1)$ such that

$$y = \lambda a + (1 - \lambda)x \Rightarrow y - a = (\lambda - 1)a + (1 - \lambda)x$$

$$\Rightarrow y - a = (1 - \lambda)(x - a)$$

$$f \text{ is convex} \Rightarrow f(y) \leq \lambda f(a) + (1 - \lambda)f(x)$$


$$\Rightarrow f(y) - f(a) \leq (\lambda - 1)f(a) + (1 - \lambda)f(x)$$

$$\Rightarrow f(y) - f(a) \leq (1 - \lambda)(f(x) - f(a))$$

$$\text{Dividing both sides by } y - a = (1 - \lambda)(x - a) > 0 \Rightarrow \frac{f(y) - f(a)}{y - a} \leq \frac{f(x) - f(a)}{x - a}$$

$$\text{If } y \rightarrow a^+, \text{ then } f'(a) \leq \frac{f(x) - f(a)}{x - a} \Rightarrow f(x) \geq f(a) + f'(a)(x - a) \text{ if } x, a \in I.$$

If $a > x$ then the proof is similar and if $a = x$ then the statement is obvious.




Proof of (2) \Rightarrow (3): Let $T_a(x) = f(a) + f'(a)(x - a)$.

$$\text{If } a, b \in I, a < b \Rightarrow T_a(a) = f(a) \geq T_b(a) \text{ and } T_a(b) \leq f(b) = T_b(b)$$

$$\Rightarrow f'(a) = \frac{T_a(b) - T_a(a)}{b - a} = \frac{T_a(b) - f(a)}{b - a} \leq \frac{f(b) - T_b(a)}{b - a} = \frac{T_b(b) - T_b(a)}{b - a} = f'(b)$$

$\Rightarrow f'$ is monotonically increasing on I



Proof of (3) \Rightarrow (1): Let $a, b \in I, a < b, \lambda \in (0, 1)$ for which $x = \lambda a + (1 - \lambda)b$

$$\Rightarrow x - a = (1 - \lambda)(b - a)$$

$$b - x = \lambda(b - a)$$

Then by Lagrange's mean value theorem there exist $c_1 \in (a, x)$ and $c_2 \in (x, b)$ such that

$$\frac{f(x) - f(a)}{x - a} = f'(c_1) \text{ and } f'(c_2) = \frac{f(b) - f(x)}{b - x}.$$

f' is monotonically increasing $\Rightarrow f'(c_1) \leq f'(c_2)$

$$\Rightarrow \frac{f(x) - f(a)}{x - a} \leq \frac{f(b) - f(x)}{b - x} \Rightarrow \frac{f(x) - f(a)}{(1 - \lambda)(b - a)} \leq \frac{f(b) - f(x)}{\lambda(b - a)} \Rightarrow f(x) \leq \lambda f(a) + (1 - \lambda)f(b)$$

$\Rightarrow f$ is convex on I .

Consequence (Necessary and sufficient condition for convexity).

Assume that f is twice differentiable on the interval I . Then

(1) $f''(x) \geq 0 \forall x \in I$ if and only if f is convex on I .

(2) $f''(x) \leq 0 \forall x \in I$ if and only if f is concave on I .

Consequence.


Assume that f is twice differentiable on the interval I . Then

(1) If $f''(x) > 0 \forall x \in I$ then f is strictly convex on I .

(2) If $f''(x) < 0 \forall x \in I$ then f is strictly concave on I .

Inflection point

Definition. Assume that f is continuous at $a \in \text{int } D_f$ and there exists $\delta > 0$ such that f is convex on $(a - \delta, a)$ and concave on $(a, a + \delta)$ or f is concave on $(a - \delta, a)$ and convex on $(a, a + \delta)$. Then a is called a point of inflection of the function f .



Theorem (Necessary condition for an inflection point, second derivative test).

If f is twice differentiable at x_0 and f has an inflection point at x_0 then $f''(x_0) = 0$.

Proof. If f is convex on $(x_0 - \delta, x_0]$ and concave on $[x_0, x_0 + \delta)$ then f' is monotonically increasing on $(x_0 - \delta, x_0]$ and monotonically decreasing on $[x_0, x_0 + \delta)$
 $\Rightarrow f'$ has a local maximum at $x_0 \Rightarrow f''(x_0) = 0$.

Theorem (Sufficient condition for an inflection point, second derivative test).

If f is twice differentiable in a neighbourhood of x_0 ,
 $f''(x_0) = 0$ and f'' changes sign at x_0 ,
 then f has an inflection point at x_0 .

Theorem (Sufficient condition for an inflection point, third derivative test).

If f is three times differentiable in a neighbourhood of x_0 ,
 $f''(x_0) = 0$ and $f'''(x_0) \neq 0$,
 then f has an inflection point at x_0 .

Inflection point and higher order derivatives

Theorem. (1) Assume that f is $2k + 1$ times differentiable at x_0 , $k \geq 1$.

If $f''(x_0) = \dots = f^{(2k)}(x_0) = 0$ and $f^{(2k+1)}(x_0) \neq 0$
 then f has an inflection point at x_0 .

(2) Assume that f is $2k$ times differentiable at x_0 , $k \geq 1$.

If $f''(x_0) = \dots = f^{(2k-1)}(x_0) = 0$ and $f^{(2k)}(x_0) \neq 0$, then f is strictly convex or concave in a neighbourhood of x_0 , so f doesn't have an inflection point at x_0 .

Remark. Part (1) in other words: If the first non-zero derivative (after the second one) has an odd order then f has a local extremum at x_0 .

Linear asymptotes

Definition. The straight line $x = a$ is a **vertical asymptote** of the function f if

$$\lim_{x \rightarrow a^+} f(x) = \pm\infty \text{ or } \lim_{x \rightarrow a^-} f(x) = \pm\infty.$$

Definition. The straight line $g(x) = Ax + B$ is a **linear asymptote** of the function f at ∞ or $-\infty$ if

$$\lim_{x \rightarrow \infty} (f(x) - g(x)) = 0 \text{ or } \lim_{x \rightarrow -\infty} (f(x) - g(x)) = 0.$$



$g(x)$ is a **horizontal asymptote** if $A = 0$ and an **oblique or slant asymptote** if $A \neq 0$.

Statement. $g(x) = Ax + B$ is a linear asymptote of f at $\pm\infty$ if and only if

$$A = \lim_{x \rightarrow \pm\infty} \frac{f(x)}{x} \text{ and } B = \lim_{x \rightarrow \pm\infty} (f(x) - Ax)$$

Example. $\lim_{x \rightarrow \frac{\pi}{2} \pm} \tan x = \mp\infty \implies x = \frac{\pi}{2}$ is a vertical asymptote of $f(x) = \tan(x)$.

Example. If $f(x) = x + 2 + \frac{1}{x}$ then $g(x) = x + 2$ is a linear asymptote of f at $\pm\infty$.



Example. If $f(x) = x e^{\frac{2}{x}}$ then $g(x) = x + 2$ is a linear asymptote of f at $\pm\infty$.

Solution. $A = \lim_{x \rightarrow \pm\infty} \frac{f(x)}{x} = \lim_{x \rightarrow \pm\infty} \frac{x e^{\frac{2}{x}}}{x} = \lim_{x \rightarrow \pm\infty} e^{\frac{2}{x}} = e^0 = 1$

$$B = \lim_{x \rightarrow \pm\infty} \left(x e^{\frac{2}{x}} - x \right) = \lim_{x \rightarrow \pm\infty} \frac{e^{\frac{2}{x}} - 1}{\frac{1}{x}}. \text{ Let } y = \frac{2}{x}, \text{ then } B = \lim_{y \rightarrow 0 \pm} \frac{e^y - 1}{\frac{1}{2} \cdot y} = 2,$$

using that $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$. The limit can also be calculate with the L'Hospital's rule.

So $g(x) = x + 2$.

Extreme values on a closed interval

Remark. If f is continuous on a closed and bounded interval then by the Weierstrass extreme value theorem f has a minimum and a maximum.

The possible points are:

- 1) the points where f is not differentiable
- 2) the points where the derivative of f is 0
- 3) the endpoints of the interval

Finally the largest and smallest of the possible values must be selected.

Analyzing graphs of functions

Summary of the steps:

- 1) finding the domain of f
- 2) finding the zeros of f
- 3) parity, periodicity
- 4) limits at the endpoints of the intervals constituting the domain
- 5) investigation of f' \implies monotonicity, extreme values
- 6) investigation of f'' \implies convexity/concavity, inflection points
- 7) linear asymptotes
- 8) plotting the graph of f , finding the range of f

Exercises

<https://math.bme.hu/~nagyi/calculus1/functions.pdf>

Examples

$$1. f(x) = \frac{x}{x^3 + 1}$$

$$D_f = (-\infty, -1) \cup (-1, \infty); f(x) = 0 \iff x = 0;$$

$$\lim_{x \rightarrow \pm\infty} f(x) = 0, \quad \lim_{x \rightarrow -1+0} f(x) = -\infty, \quad \lim_{x \rightarrow -1-0} f(x) = +\infty$$

Monotonicity, local extremum:

$$f'(x) = \frac{1 - 2x^3}{(x^3 + 1)^2} = 0 \iff x = \frac{1}{\sqrt[3]{2}} \approx 0.79$$


x	$x < -1$	$-1 < x < \frac{1}{\sqrt[3]{2}}$	$x = \frac{1}{\sqrt[3]{2}}$	$x > \frac{1}{\sqrt[3]{2}}$
f'	+	+	0	-
f	\nearrow	\nearrow	max: $\frac{\sqrt[3]{4}}{3} \approx 0.53$	\searrow

Convexity / concavity, inflection points:

$$f''(x) = \frac{6x^2(x^3 - 2)}{(x^3 + 1)^3} = 0 \iff x = 0 \text{ or } x = \sqrt[3]{2} \approx 1.26$$

x	$x < -1$	$-1 < x < 0$	$x = 0$	$0 < x < \sqrt[3]{2}$	$x = \sqrt[3]{2}$	$x > \sqrt[3]{2}$
f''	+	-	0	-	0	+
f	\cup	\cap		\cap	infl: $\frac{\sqrt[3]{2}}{3} \approx 0.42$	\cup

The graph of f :



2. $f(x) = 2 \sin x + \sin 2x$

$D_f = \mathbb{R}$; f is odd;

f is periodic with period $2\pi \Rightarrow$ it may be assumed that $0 \leq x \leq 2\pi$;

\Rightarrow on this interval $f(x) = 0 \Leftrightarrow x = 0$ or $x = \pi$ or $x = 2\pi$

Monotonicity, local extremum:

$$f'(x) = 2 \cos x + 2 \cos 2x = 2(\cos x + \cos^2 x - (1 - \cos^2 x)) =$$

$$= 2 \cdot (2 \cos^2 x + \cos x - 1) = 0 \Rightarrow (\cos x)_{1,2} = \frac{-1 \pm 3}{4} \Rightarrow \cos x = -1 \text{ or } \cos x = \frac{1}{2}$$

$$\Rightarrow x_1 = \frac{\pi}{3}, x_2 = \pi, x_3 = \frac{5\pi}{3}$$

x	\emptyset	$(\emptyset, \frac{\pi}{3})$	$\frac{\pi}{3}$	$(\frac{\pi}{3}, \pi)$	π	$(\pi, \frac{5\pi}{3})$	$\frac{5\pi}{3}$	$(\frac{5\pi}{3}, 2\pi)$	2π
f'	+	+	\emptyset	-	\emptyset	-	\emptyset	+	+
f		\nearrow	$\max: \frac{3\sqrt{3}}{2}$	\searrow		\searrow	$\min: -\frac{3\sqrt{3}}{2}$	\nearrow	

Convexity / concavity, inflection points:


$$f''(x) = -2 \sin x - 4 \sin 2x = -2 \sin x - 8 \sin x \cos x =$$

$$= -2 \sin x(1 + 4 \cos x) = 0 \Rightarrow \sin x = 0 \text{ or } \cos x = -\frac{1}{4}$$

$$\Rightarrow x_1 = 0, x_2 = \pi, x_3 = 2\pi, x_4 = \arccos\left(-\frac{1}{4}\right) \approx 1.82, x_5 = 2\pi - \arccos\left(-\frac{1}{4}\right) \approx 4.46$$

x	\emptyset	$(\emptyset, 1.82)$	1.82	$(1.82, \pi)$	π	$(\pi, 4.46)$	4.46	$(4.46, 2\pi)$	2π
f''	\emptyset	-	\emptyset	+	\emptyset	-	\emptyset	+	\emptyset
f	infl: \emptyset	\cap	infl: \searrow $\frac{3\sqrt{15}}{8}$	\cup	infl: \emptyset	\cap	infl: \searrow $-\frac{3\sqrt{15}}{8}$	\cup	infl: \emptyset

The graph of f :



Implicitly given curve

Example. The curve $y = y(x)$ is given by the following implicit equation:

$$x \sinh x - y \cosh y = 0$$

Study the properties of this curve in a neighbourhood of $(0, 0)$.

Solution. The point $(0, 0)$ is on the curve: $y(0) = 0$.

1) The first derivative of $x \sinh x - y(x) \cosh y(x) = 0$ with respect to x :

$$\sinh x + x \cosh x - y'(x) \cosh y(x) - y(x) y'(x) \sinh y(x) = 0$$

$$\text{If } x = 0, y = 0 \implies 0 + 0 \cdot 1 - y'(0) \cdot 1 - 0 \cdot y'(0) \cdot 0 = 0 \implies y'(0) = 0$$

2) The second derivative with respect to x :

$$\cosh x + \cosh x + x \sinh x - y''(x) \cosh y(x) - y'(x) y'(x) \sinh y(x) - y'(x) y'(x) \sinh y(x) - y(x) y''(x) \sinh y(x) - y(x) y'(x) y'(x) \cosh y(x) = 0$$

$$\text{If } x = 0, y = 0 \implies 1 + 1 + 0 - y''(0) - 0 - 0 - 0 - 0 = 0 \implies y''(0) = 2$$

Since $y'(0) = 0$ and $y''(0) = 2 > 0$ then the curve $y = y(x)$ has local minimum at $x = 0$ and it is convex in some neighbourhood of $x = 0$.

