Exercises

Analyze the following functions and sketch their graphs.

1. $f(x) = x^4 - 4x^3$
2. $f(x) = -x^4 + 18x^2$
3. $f(x) = x^5 + 5x^4$
4. $f(x) = \dfrac{1}{1+x^2}$
5. $f(x) = \dfrac{1}{1+x^2}$
6. $f(x) = \dfrac{x}{1+x^2}$
7. $f(x) = \dfrac{x}{1-x^2}$
8. $f(x) = \dfrac{x}{(1-2x)^2}$
9. $f(x) = \dfrac{x^2}{x+1}$
10. $f(x) = \dfrac{x^3}{x^2-3}$
11. $f(x) = x\,e^{-x}$
12. $f(x) = (x+2)^2\,e^{-x}$
13. $f(x) = e^{-x^2}$
14. $f(x) = x\,e^{-x^2}$
15. $f(x) = x^2 \ln x$
16. $f(x) = \arctan(x^2)$

Summary of the steps:

1) finding the domain of f
2) finding the zeros of f
3) parity, periodicity
4) limits at the endpoints of the intervals constituting the domain
5) investigation of f' $\Longrightarrow$ monotonicity, extreme values
6) investigation of f'' $\Longrightarrow$ convexity/concavity, inflection points
7) linear asymptotes
8) plotting the graph of f, finding the range of f

Theorems:

1) Assume that f is differentiable at $x_0 \in \operatorname{int} D_f$.

Necessary condition for a local extremum at x_0:

$f'(x_0) = 0$

Sufficient condition for a local extremum at x_0:

a) $f'(x_0) = 0$ and f' changes sign at x_0

b) f is twice differentiable at x_0, $f'(x_0) = 0$ and $f''(x_0) \neq 0$

($f''(x_0) > 0$: local minimum, $f''(x_0) < 0$: local maximum)

2) Assume that f is twice differentiable at $x_0 \in \operatorname{int} D_f$.

Necessary condition for an inflection point at x_0:

$f''(x_0) = 0$

Sufficient condition for an inflection point at x_0:

a) $f''(x_0) = 0$ and f'' changes sign at x_0

b) f is three times differentiable at x_0, $f''(x_0) = 0$ and $f'''(x_0) \neq 0$

Asymptotes:

1) The straight line $x = a$ is a vertical asymptote of the function f if $\lim\limits_{x\to a+} f(x) = \pm\infty$ or $\lim\limits_{x\to a-} f(x) = \pm\infty$.

2) The straight line $g(x) = A\,x + B$ is a linear asymptote of the function f at ∞ or $-\infty$ if

$\lim\limits_{x\to\infty} (f(x) - g(x)) = 0$ or $\lim\limits_{x\to-\infty} (f(x) - g(x)) = 0$. Then $A = \lim\limits_{x\to\pm\infty} \dfrac{f(x)}{x}$ and $B = \lim\limits_{x\to\pm\infty} (f(x) - A\,x)$.

Solutions

1. $f(x) = x^4 - 4x^3$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$ or $x = 4$

$f'(x) = 4x^3 - 12x^2 = 4x^2(x-3) = 0 \iff x = 0$ or $x = 3$

$f''(x) = 12x^2 - 24x = 12x(x-2) = 0 \iff x = 0$ or $x = 2$

$\lim_{x\to-\infty} f(x) = \lim_{x\to+\infty} f(x) = +\infty$

$R_f = [-27, \infty)$

x	x<0	x=0	0<x<3	x=3	x>3
f'	−	0	−	0	+
f	↘		↘	min:−27	↗

x	x<0	x=0	0<x<2	x=2	x>2
f''	+	0	−	0	+
f	∪	infl:0	∩	infl:−16	∪



2. $f(x) = -x^4 + 18x^2$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$ or $x = \pm 3\sqrt{2}$; f is even;

$f'(x) = -4x^3 + 36x = 4x(-x^2 + 9) = 0 \iff x = 0$ or $x = \pm 3$

$f''(x) = -12x^2 + 36 = 0 \iff x = \pm\sqrt{3}$

$\lim_{x\to-\infty} f(x) = \lim_{x\to+\infty} f(x) = -\infty$

$R_f = (-\infty, 81]$

x	x<−3	x=−3	−3<x<0	x=0	0<x<3	x=3	x>3
f'	+	0	−	0	+	0	−
f	↗	max:81	↘	min:0	↗	max:81	↘

x	$x<-\sqrt{3}$	$x=-\sqrt{3}$	$-\sqrt{3}<x<\sqrt{3}$	$x=\sqrt{3}$	$x>\sqrt{3}$
f''	−	0	+	0	−
f	∩	infl:45	∪	infl:45	∩



3. $f(x) = x^5 + 5x^4$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$ or $x = -5$

$f'(x) = 5x^4 + 20x^3 = 5x^3(x+4) = 0 \iff x = -4$ or $x = 0$

$f''(x) = 20x^3 + 60x^2 = 20x^2(x+3) = 0 \iff x = -3$ or $x = 0$

$\lim_{x\to-\infty} f(x) = -\infty$, $\lim_{x\to+\infty} f(x) = +\infty$

$R_f = \mathbb{R}$

x	x<−4	x=−4	−4<x<0	x=0	x>0
f'	+	0	−	0	+
f	↗	max:256	↘	min:0	↗

x	x<−3	x=−3	−3<x<0	x=0	x>0
f''	−	0	+	0	+
f	∩	infl:162	∪		∪



4. $f(x) = \dfrac{1}{1+x^2}$

$D_f = \mathbb{R}$; $f(x) \neq 0$; f is even

$f'(x) = -\dfrac{2x}{(1+x^2)^2} = 0 \iff x = 0$

$f''(x) = \dfrac{2\cdot(-1+3x^2)}{(1+x^2)^3} = 0 \iff x = \pm\dfrac{1}{\sqrt{3}}$

$\lim_{x\to-\infty} f(x) = 0$, $\lim_{x\to+\infty} f(x) = 0$

horizontal asymptote: $y = 0$

$R_f = (0, 1]$

x	x<0	x=0	x>0
f'	+	0	−
f	↗	max:1	↘

x	$x<-\frac{1}{\sqrt{3}}$	$x=-\frac{1}{\sqrt{3}}$	$-\frac{1}{\sqrt{3}}<x<\frac{1}{\sqrt{3}}$	$x=\frac{1}{\sqrt{3}}$	$x>\frac{1}{\sqrt{3}}$
f''	+	0	−	0	+
f	∪	infl: $\frac{3}{4}$	∩	infl: $\frac{3}{4}$	∪



5. $f(x) = \dfrac{1}{1 - x^2}$

$D_f = \mathbb{R} \setminus \{-1, 1\}$; $f(x) \neq 0$; f is even

$$f'(x) = \frac{2x}{(1 - x^2)^2} = 0 \iff x = 0$$

$$f''(x) = \frac{2 \cdot (1 + 3x^2)}{(1 - x^2)^3} \neq 0$$

$$\lim_{x \to \pm\infty} f(x) = 0, \quad \lim_{x \to -1 \pm 0} f(x) = \pm\infty, \quad \lim_{x \to 1 \pm 0} f(x) = \mp\infty$$

horizontal asymptote : $y = 0$

vertical asymptotes: $x = 1$, $x = -1$

$R_f = (-\infty, 0) \cup [1, +\infty)$

x	x<−1	−1<x<0	x=0	0<x<1	1<x
f'	−	−	0	+	+
f	↘	↘	min:1	↗	↗

x	x<−1	−1<x<1	x>1
f''	−	+	−
f	∩	∪	∩



6. $f(x) = \dfrac{x}{1 + x^2}$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$; f is odd

$$f'(x) = \frac{1 - x^2}{(1 + x^2)^2} = 0 \iff x = \pm 1$$

$$f''(x) = \frac{2x(-3 + x^2)}{(1 + x^2)^3} = 0 \iff x = 0 \text{ or } x = \pm\sqrt{3}$$

$$\lim_{x \to -\infty} f(x) = 0, \quad \lim_{x \to +\infty} f(x) = 0$$

horizontal asymptote : $y = 0$

$R_f = \left[-\frac{1}{2}, \frac{1}{2}\right]$

x	x<−1	x=−1	−1<x<1	x=1	x>1
f'	−	0	+	0	−
f	↘	min: $-\frac{1}{2}$	↗	max: $\frac{1}{2}$	↘



x	x<$-\sqrt{3}$	x=$-\sqrt{3}$	$-\sqrt{3}$<x<0	x=0	0<x<$\sqrt{3}$	x=$\sqrt{3}$	x>$\sqrt{3}$
f''	−	0	+	0	−	0	+
f	∩	infl: $-\frac{\sqrt{3}}{4}$	∪	infl:0	∩	infl: $\frac{\sqrt{3}}{4}$	∪

7. $f(x) = \dfrac{x}{1 - x^2}$

$D_f = \mathbb{R} \setminus \{-1, 1\}$

$f(x) = 0 \iff x = 0;\ f$ is odd

$f'(x) = \dfrac{1 + x^2}{(1 - x^2)^2} \neq 0$

$f''(x) = \dfrac{2x(3 + x^2)}{(1 - x^2)^3} = 0 \iff x = 0$

$\lim\limits_{x \to \pm\infty} f(x) = 0, \quad \lim\limits_{x \to -1 \pm 0} f(x) = \mp\infty, \quad \lim\limits_{x \to 1 \pm 0} f(x) = \mp\infty$

horizontal asymptote : $y = 0$

vertical asymptotes: $x = 1,\ x = -1$

$R_f = \mathbb{R}$



x	x<−1	−1<x<1	x>1
f'	+	+	+
f	↗	↗	↗

x	x<−1	−1<x<0	x=0	0<x<1	1<x
f''	+	−	0	+	−
f	∪	∩	infl:0	∪	∩

8. $f(x) = \dfrac{x}{(1 - 2x)^2}$

$D_f = \mathbb{R} \setminus \left\{\dfrac{1}{2}\right\}$

$f(x) = 0 \iff x = 0$

$f'(x) = \dfrac{1 + 2x}{(1 - 2x)^3} = 0 \iff x = -\dfrac{1}{2}$

$f''(x) = \dfrac{8 \cdot (1 + x)}{(1 - 2x)^4} = 0 \iff x = -1$

$\lim\limits_{x \to \pm\infty} f(x) = 0, \quad \lim\limits_{x \to -\frac{1}{2} \pm 0} f(x) = +\infty$

horizontal asymptote : $y = 0$

vertical asymptote: $x = \dfrac{1}{2}$

$R_f = \left[-\dfrac{1}{8}, +\infty\right)$



x	$x<-\frac{1}{2}$	$x=-\frac{1}{2}$	$-\frac{1}{2}<x<\frac{1}{2}$	$x>\frac{1}{2}$
f'	−	0	+	−
f	↘	min: $-\frac{1}{8}$	↗	↘

x	x<−1	x=−1	$-1<x<\frac{1}{2}$	$x>\frac{1}{2}$
f''	−	0	+	−
f	∩	infl: $-\frac{1}{9}$	∪	∩

9. $f(x) = \dfrac{x^2}{x+1}$

$D_f = \mathbb{R} \setminus \{-1\}$;

$f(x) = 0 \iff x = 0$

$f'(x) = \dfrac{x(2+x)}{(1+x)^2} = 0 \iff x = -2 \text{ or } x = 0$

$f''(x) = \dfrac{2}{(1+x)^3} \neq 0$

$\lim\limits_{x\to\pm\infty} f(x) = \pm\infty, \quad \lim\limits_{x\to -1\pm 0} f(x) = \pm\infty$

vertical asymptote : $x = -1$

linear asymptote: $y = x - 1$

$R_f = \mathbb{R}$

x	x<−2	x=−2	−2<x<−1	−1<x<0	x=0	x>0
f'	+	0	−	−	0	+
f	↗	max:−4	↘	↘	min:0	↗



x	x<−1	x>−1
f''	−	+
f	∩	∪

10. $f(x) = \dfrac{x^3}{x^2-3}$

$D_f = \mathbb{R} \setminus \left\{-\sqrt{3}, \sqrt{3}\right\}$;

$f(x) = 0 \iff x = 0$; f is odd

$f'(x) = \dfrac{x^2(x^2-9)}{(x^2-3)^2} = 0 \iff x = 0 \text{ or } x = \pm 3$

$f''(x) = \dfrac{6x(9+x^2)}{(x^2-3)^3} = 0 \iff x = 0$

$\lim\limits_{x\to\pm\infty} f(x) = \pm\infty, \quad \lim\limits_{x\to -\sqrt{3}\pm 0} f(x) = \pm\infty, \quad \lim\limits_{x\to \sqrt{3}\pm 0} f(x) = \pm\infty$

vertical asymptotes : $x = -\sqrt{3}$, $x = \sqrt{3}$

linear asymptote: $y = x$

$R_f = \mathbb{R}$

x	x<−3	x=−3	$-3<x<-\sqrt{3}$	$-\sqrt{3}<x<0$	x=0	$0<x<\sqrt{3}$	$\sqrt{3}<x<3$	x=3	x>3
f'	+	0	−	−	0	−	−	0	+
f	↗	max: $-\frac{9}{2}$	↘	↘		↘	↘	min: $\frac{9}{2}$	↗

x	$x<-\sqrt{3}$	$-\sqrt{3}<x<0$	x=0	$0<x<\sqrt{3}$	$x>\sqrt{3}$
f''	−	+	0	−	+
f	∩	∪	infl:0	∩	∪



11. $f(x) = x\,e^{-x}$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$

$f'(x) = e^{-x}(1 - x) = 0 \iff x = 1$

$f''(x) = e^{-x}(x - 2) = 0 \iff x = 2$

$\lim\limits_{x\to+\infty} f(x) = \lim\limits_{x\to+\infty} \frac{x}{e^x} \overset{L'H}{=} \lim\limits_{x\to+\infty} \frac{1}{e^x} = 0$

$\lim\limits_{x\to-\infty} f(x) = -\infty$

$R_f = \left(-\infty, \frac{1}{e}\right]$

x	x<1	x=1	x>1
f'	+	0	−
f	↗	max: $\frac{1}{e}$ ≈0.37	↘

x	x<2	x=2	x>2
f''	−	0	+
f	⋂	infl: $\frac{2}{e^2}$ ≈0.27	⋃



12. $f(x) = (x+2)^2\,e^{-x}$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = -2$

$f'(x) = -e^{-x}\,x\,(2 + x) = 0 \iff x = 0$ or $x = -2$

$f''(x) = e^{-x}\left(x^2 - 2\right) = 0 \iff x = \pm\sqrt{2}$

$\lim\limits_{x\to+\infty} f(x) = \lim\limits_{x\to+\infty} \frac{(x+2)^2}{e^x} \overset{L'H}{=} \lim\limits_{x\to+\infty} \frac{2\,(x+2)}{e^x} \overset{L'H}{=} \lim\limits_{x\to+\infty} \frac{2}{e^x} = 0$

$\lim\limits_{x\to-\infty} f(x) = +\infty$

$R_f = [0, +\infty)$

x	x<−2	x=−2	−2<x<0	x=0	x>0
f'	−	0	+	0	−
f	↘	min:0	↗	max:4	↘



x	x<−$\sqrt{2}$	x=−$\sqrt{2}$	−$\sqrt{2}$<x<$\sqrt{2}$	x=$\sqrt{2}$	x>$\sqrt{2}$
f''	+	0	−	0	+
f	⋃	infl:≈1.41	⋂	infl:≈2.83	⋃

13. $f(x) = e^{-x^2}$

$D_f = \mathbb{R}$; $f(x) \neq 0$; f is even

$f'(x) = e^{-x^2}(-2x) = 0 \iff x = 0$

$f''(x) = 2e^{-x^2}(-1 + 2x^2) = 0 \iff x = \pm\frac{1}{\sqrt{2}}$

$\lim_{x\to\pm\infty} f(x) = 0$; $R_f = (0, 1]$

x	x<0	x=0	x>0
f'	+	0	−
f	↗	max:1	↘

x	$x<-\frac{1}{\sqrt{2}}$	$x=-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}<x<\frac{1}{\sqrt{2}}$	$x=\frac{1}{\sqrt{2}}$	$x>\frac{1}{\sqrt{2}}$
f''	+	0	−	0	+
f	∪	infl: $\frac{1}{\sqrt{e}}\approx 0.61$	∩	infl: $\approx\frac{1}{\sqrt{e}}\approx 0.61$	∪





14. $f(x) = x\,e^{-x^2}$

$D_f = \mathbb{R}$; $f(x) = 0 \iff x = 0$; f is odd

$f'(x) = -e^{-x^2}(-1 + 2x^2) = 0 \iff x = \pm\frac{1}{\sqrt{2}}$

$f''(x) = 2e^{-x^2}x(-3 + 2x^2) = 0 \iff x = 0$ or $x = \pm\sqrt{\frac{3}{2}}$

$\lim_{x\to\pm\infty} f(x) = \lim_{x\to\pm\infty}\frac{x}{e^{x^2}} \overset{L'H}{=} \lim_{x\to\pm\infty}\frac{1}{e^{x^2}\cdot 2x} = 0$

$R_f = \left[-\frac{1}{\sqrt{2e}}, \frac{1}{\sqrt{2e}}\right]$

x	$x<-\frac{1}{\sqrt{2}}$	$x=-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}<x<\frac{1}{\sqrt{2}}$	$x=\frac{1}{\sqrt{2}}$	$x>\frac{1}{\sqrt{2}}$
f'	−	0	+	0	−
f	↘	min: $-\frac{1}{\sqrt{2e}}\approx -0.43$	↗	max: $\frac{1}{\sqrt{2e}}\approx 0.43$	↘

x	$x<-\sqrt{\frac{3}{2}}$	$x=-\sqrt{\frac{3}{2}}$	$-\sqrt{\frac{3}{2}}<x<0$	x=0	$0<x<\sqrt{\frac{3}{2}}$	$x=\sqrt{\frac{3}{2}}$	$x>\sqrt{\frac{3}{2}}$
f''	−	0	+	0	−	0	+
f	∩	infl: ≈ -0.27	∪	infl:0	∩	infl: ≈ 0.27	∪

15. $f(x) = x^2 \ln x$

$D_f = \mathbb{R}^+;\ f(x) = 0 \iff x = 1$

$f'(x) = x(1 + 2\ln x) = 0 \iff x = \frac{1}{\sqrt{e}} \approx 0.61$

$f''(x) = 3 + 2\ln x = 0 \iff x = \frac{1}{e^{3/2}} \approx 0.22$

$\lim_{x\to 0+0} f(x) = \lim_{x\to 0+0} \frac{\ln x}{\frac{1}{x^2}} \overset{L'H}{=} \lim_{x\to 0+0} \frac{\frac{1}{x}}{-\frac{2}{x^3}} = \lim_{x\to 0+0} \frac{-x^2}{2} = 0$

$\lim_{x\to +\infty} f(x) = +\infty$

$R_f = \left[-\frac{1}{2e}, +\infty\right)$

x	$0<x<\frac{1}{\sqrt{e}}$	$x=\frac{1}{\sqrt{e}}$	$x>\frac{1}{\sqrt{e}}$
f'	−	0	+
f	↗	min: $-\frac{1}{2e} \approx -0.18$	↘

x	$0<x<\frac{1}{e^{3/2}}$	$x=\frac{1}{e^{3/2}}$	$x>\frac{1}{e^{3/2}}$
f''	−	0	+
f	∩	infl: $-\frac{3}{2e^3} \approx -0.07$	∪



16. $f(x) = \arctan(x^2)$

$D_f = \mathbb{R};\ f(x) = 0 \iff x = 0;\ f$ is even

$f'(x) = \frac{2x}{1+x^4} = 0 \iff x = 0$

$f''(x) = \frac{2\cdot(1-3x^4)}{(1+x^4)^2} = 0 \iff x = \pm\frac{1}{\sqrt[4]{3}}$

$\lim_{x\to\pm\infty} f(x) = \frac{\pi}{2};\quad R_f = \left[0, \frac{\pi}{2}\right)$

x	x<0	0	x>0
f'	−	0	+
f	↘	min:0	↗



x	$x<-\frac{1}{\sqrt[4]{3}}$	$x=-\frac{1}{\sqrt[4]{3}}$	$-\frac{1}{\sqrt[4]{3}}<x<\frac{1}{\sqrt[4]{3}}$	$x=\frac{1}{\sqrt[4]{3}}$	$x>\frac{1}{\sqrt[4]{3}}$
f''	−	0	+	0	−
f	∩	infl: $\frac{\pi}{6}$	∪	infl: $\frac{\pi}{6}$	∩